

- **The Association between Infant Feeding Patterns and Type-I Diabetes in Children at Ministry of National Guard, Health Affairs Hospital, Riyadh.**

Reem F Alshammari

- **Prevalence and Antibiotic Resistance of Cronobacter Spp. Associated with Powdered Infant Milk Formulas and Dried Milks in Saudi Arabia.**

Khalid. M. Aljarallah

- **Athletic Pubelgia-A Review of Literature.**

Qassim I. Muaidi





# MJHS

## MAJMAAH JOURNAL OF HEALTH SCIENCES

A JOURNAL PUBLISHED BY MAJMAAH UNIVERSITY

### Editorial Board

#### EDITOR-IN-CHIEF

#### **Prof. Mohammed AlRukban**

Professor of Family Medicine, College of Medicine,  
King Saud University, Riyadh

#### EDITORS

##### **Prof. Abdulrahman M. Almazrou**

Department of Pediatrics, KSU Consultant, Pediatrics  
Infectious Diseases Chief Supervisor & Dean AlRajhi  
colleges, Qassim, KSA

##### **Prof. Farooq Khan**

Professor of Medicine, State University of New York.  
Director Research and Publication  
Stony Brook, New York

##### **Prof. Omar Hasan K Kasule**

Faculty of Medicine at King Fahd Medical City,  
Riyadh, KSA

##### **Prof. Mohammad Faheem**

Head Department of Physiology, HIMSAR, Jamia  
Hamdard, New Delhi, India.

##### **Dr. Ibrahim Alhoqail**

Associate Professor Dermatology, College of  
Medicine, King Saud University

##### **Dr. Abdullah Ali Alghasham**

Associate Professor, College of Medicine, Qaseem  
University, KSA

##### **Prof. Samuel Lee**

Professor of Medicine, University of Calgary, Canada

#### EDITORIAL ASSISTANTS

##### **Dr. Khalid Tohami**

Assistant Professor, Community Medicine,  
Department of Public  
Health & Community Medicine,  
College of Medicine, Majmaah University

##### **Dr. Moattar Raza Rizvi**

Assistant Professor, Physiology  
Department of Nursing,  
College of Applied Medical Sciences,  
Majmaah University.

##### **Dr. Fuzail Ahmad**

Head Department of Physical Therapy & Health  
Rehabilitation,  
College of Applied Medical Sciences,  
Majmaah University

##### **Mr. Waqas Sami**

Lecturer, Biostatistics, Department of  
Public Health & Community Medicine,  
College of Medicine, Majmaah University

# MJHS

## MAJMAAH JOURNAL OF HEALTH SCIENCES

A JOURNAL PUBLISHED BY MAJMAAH UNIVERSITY

---

### TABLE OF CONTENTS

	Page No.
<b>Preface</b>	
<b>ORIGINAL ARTICLES</b>	
1. The Association between Infant Feeding Patterns and Type-I Diabetes In Children At Ministry Of National Guard, Health Affairs Hospital, Riyadh. <i>Reem F AlShammari</i>	1-12
2. Prevalence and Antibiotic Resistance of Cronobacter Spp. Associated with Powdered Infant Milk Formulas and Dried Milks in Saudi Arabia. <i>Khaled M. AlJarallah</i>	13-18
3. Fast Food Consumption among University Students, Saudi Arabia. A Cross-Sectional Study. <i>Fahad AlFahied, Waqas Sami, Yasser AlTowyan, Turki AlJohani, Abdulaziz AlHaddad, Abdullah AlKhamisi</i>	19-24
<b>Review Article</b>	
4. Athletic Pubalgia – A Review of Literature. <i>Qassim I. Muaidi</i>	25-35
<b>VIEW POINT</b>	
5. Teaching biostatistics to medical undergraduates in integrated curriculum. Viewpoint. <i>Waqas Sami, Dr. Mohammad AlMansour, Tayyaba Waqas</i>	36-39
<b>CASE REPORT</b>	
6. Chondroid Lipoma: A Rare Recently Described Benign Lipomatous Tumor. <i>Salman T. AlMalki, Abdullah S. AlKhamisi</i>	40-44
<b>Medical News</b>	45-47
<b>Medical Quiz</b>	48
<b>Upcoming Conferences</b>	49-50
<b>Publication Guidelines</b>	51-57

## PREFACE

We would like to present, with great pleasure another volume of a Majmaah Journal of Health Science MJHS. It is a great challenge to bring a new journal into the world, especially when the journal aims to publish high quality manuscripts. The main object of MJHS is to publish the research papers well in time but with peer review by subject experts. This is the 3<sup>rd</sup> year from the time we published the inaugural volume of this journal. The journal has Editorial Board of Scientist of International repute. With our editorial board's cumulative experience, this journal brings a substantial representation of the field of health sciences. Without the service and dedication of our editorial board, MJHS would have never existed. This is an open access journal which means that all content is freely available without charge to the user or his/her institution. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles in this journal without asking prior permission from the publisher or the author.

The Journal is currently in the process of getting indexed into several of databases. The success of our journal depends directly on the number of quality articles submitted for review. Accordingly, I would like to request your participation by submitting quality manuscripts for review and encouraging your colleagues to submit quality manuscripts for review. One of the great benefits we can provide to our prospective authors, regardless of acceptance of their manuscript or not, is the mentoring nature of our review process. MJHS provides authors with high quality,

helpful reviews that are shaped to assist authors in improving their manuscripts.

We thank our editors for sharing their invaluable editorial experience with us. The editorial board of MJHS has done a tremendous job; I thoroughly enjoyed the professionalism and enthusiasm of our editorial team. The journal would not be here before you without the continuous efforts of Dr. Khaled Al Tohami, Dr. Moattar Raza Rizvi, Dr. Fuzail Ahmad & Mr. Waqas Sami, the assistant to the editorial board, who kept us all on track. I thank all our reviewers, for making themselves available and providing us with timely review.

The Research papers, reviews or short communications may be sent by email to the Editor-in-Chief at the following email address: [mjhs@mu.edu.sa](mailto:mjhs@mu.edu.sa)

### Editor in Chief

**Prof. Mohammad Othman Al-Rukban**



**ORIGINAL ARTICLE**

## The Association between Infant Feeding Patterns and Type I Diabetes in Children at Ministry of National Guard, Health Affairs Hospital, Riyadh

Reem F Alshammari\*<sup>1</sup>

\* For correspondence [doctor-777@hotmail.com](mailto:doctor-777@hotmail.com)

<sup>1</sup>Department of Family Medicine and Health Care, King Abdulaziz medical city, National Guard Health affair, Riyadh, KSA

Submitted on: 3<sup>rd</sup> May 2015, Accepted on: 6<sup>th</sup> September 2015

### Abstract

**Introduction:** There is little information on the association between early infant feeding and type 1 diabetes in children in Arabian countries. This study aimed to investigate the influence of breastfeeding on the onset of diabetes in children. **Methodology:** The study included 200 subjects (99 cases and 101 controls). Their age ranged between 1 and 14 years. They attended Ministry of National Guard - Health Affairs Hospital, Riyadh, (NGHA) during the study period (September to November 2013). Mothers were interviewed by research coordinator to fill the questionnaire. Descriptive analysis of cases and controls were carried out, and Chi-square, Odds-ratios and T-test estimates were utilized for analysis. **Results.** There was no difference between the two groups with regard to intake or duration of breastfeeding, the age of initiation of formula's milk and weaning. The study showed that subjects with type 1 diabetes had higher birth weight ( $p=0.009$ ). In addition, the study showed that the current daily consumption of cow's milk in subjects with type 1 diabetes were higher significantly in patients group ( $p=0.04$ ). Those in the type 1 DM group were more likely to have a sibling with type 1 diabetes ( $p=0.005$ ). **Conclusions:** Patients did not differ significantly from the controls with regard to dietary practices.

**Key Words:** Type 1 diabetes mellitus; case-control study; infant feeding; birth weight; family history of type 1 diabetes.

### المخلص

**المقدمة:** هناك القليل من المعلومات عن العلاقة بين تغذية الرضع في وقت مبكر وداء السكري من النوع 1 في الأطفال في منطقة الخليج. **هدف الدراسة:** للتحقيق في تأثير الرضاعة الطبيعية كعامل وقائي ضد ظهور مرض السكري لدى الأطفال. **طرق البحث:** أجريت الدراسة على 200 شخصا (99 طفلاً مصاب بمرض السكري و101 طفلاً غير مصاب بالمرض) تتراوح أعمارهم ما بين 1-14 عاماً الذين حضروا لمستشفى الحرس الوطني بالرياض، المملكة العربية السعودية، خلال فترة الدراسة (من سبتمبر إلى نوفمبر 2013). وقد تم اختيار كل من الأطفال المصابين وغير مصابين بالمرض عن طريق اخذ عينات عشوائية. وأجريت مقابلات مع الأمهات لجمع البيانات عن طريق منسق الأبحاث لملء الاستبيان. **النتائج:** شملت الدراسة 200 طفلاً (101 طفلاً غير مصاب و99 طفلاً مصاب بمرض السكري). 51.2% من الأطفال الغير مصابين بالمرض كانوا من الذكور و48.8% منهم إناث و50% من الأطفال المصابين بالسكري كانوا إناث و50% من الذكور. لم يوجد فرق بين المجموعتين فيما يتعلق بتناول حليب الام أو مدة الإرضاع كما انه لم يكن هناك اختلاف كبير بين المجموعتين فيما يتعلق بسن بدء الحليب أو الفطام. السن التي بدأ فيها الطفل بأخذ الأطعمة الصلبة لأول مرة لا تختلف كثيراً بين المجموعات ( $P=0.14$ ). أيضاً أظهرت الدراسة بأن الأوزان عند الولادة للأطفال المصابين كانت اعلى من أوزان الأطفال الغير مصابين بالمرض ( $P=0.009$ ). وأظهرت نتائج الدراسة أن الاستهلاك اليومي الحالي من حليب البقر في الأطفال المصابين بمرض السكري كانت أعلى بكثير من الأطفال الغير مصابين بالمرض ( $P=0.04$ ). أيضاً مجموعة الأطفال المصابين بالمرض لديهم احتماليه أكبر ان يكون لديهم اخوه مصابين بمرض السكري ( $p=0.005$ ). **الخلاصة:** المرضى في هذه الدراسة لا يختلفون كثيراً عن الأطفال الغير مصابين بالمرض فيما يتعلق بالعادات الغذائية، أو الرضاعة الطبيعية، أو سن الأم، أو ترتيب الطفل في الأسرة. نحتاج الى دراسة تقام في المجتمع وليس في المستشفيات قد تستطيع ايضاح هذه العلاقة بشكل أفضل. في الدراسة وجد ان وزن الطفل عند الولادة، الاستهلاك اليومي لحليب الأبقار، والتاريخ العائلي لأخوة مصابين بمرض السكري من النوع 1 ترتبط بعلاقة وثيقة مع خطر الإصابة بهذا المرض. **الكلمات الدلالية:** مرض السكري من النوع 1؛ الدراسة التي تختبر المصابين وغير مصابين بالمرض؛ العوامل البيئية؛ تغذية الرضع؛ وزن الولادة؛ التاريخ العائلي لمرض السكري من النوع.

**Introduction:** Type one diabetes mellitus (T1DM) is the most common chronic disease in childhood<sup>1</sup>. The incidence of childhood type 1

disease is rising worldwide, with reported annual increase of 2-5 percent in Europe, the Middle East, and Australia.<sup>2-7</sup>

Both genetic and non-genetic factors contribute to disease risk. Nonetheless, studies of familial clustering suggest that genetics accounts for only about half of the risk fraction.<sup>8-9</sup>

The only environmental trigger undergoing active investigation is early exposure to cow's milk proteins, which may be important in T1DM pathogenesis; conversely, breast milk may protect against triggering of the autoimmunity attacks.<sup>10</sup>

A series of studies has shown that children with newly diagnosed type 1 diabetes have increased concentrations of antibodies to dietary antigens and cow milk proteins in particular.<sup>11-13</sup> It is thought to be caused by inflammation and increased gut permeability which represent an early immune aberration that predisposes to b cell autoimmunity and type 1 diabetes. Gut permeability decreases faster over the first months of life in breastfed infants compared with infants given conventional or partly hydrolyzed formulas.<sup>14</sup> Early enterovirus infections have been implicated as a strong trigger candidate for b cell autoimmunity.<sup>15</sup> It is shown that breastfeeding protects against enterovirus infections in the infant period<sup>16-17</sup> and, accordingly, this would decrease the risk of enterovirus-triggered b cell autoimmunity. Whether or not breastfeeding protects against type 1 diabetes is a controversial issue and current evidence provides contradictory results in regards to the association of breastfeeding or

early introduction of cow's milk and formula with the development of type 1 diabetes.

Although it is important to identify the association between breastfeeding and cow's milk and development of T1DM, we are not aware of any available studies in this regard in Arab countries.

This study aimed to assess the relationship between breastfeeding and development of T1DM in Saudi children.

### **Methodology**

A case - control study was conducted over the period of 3 months starting September 2013. A total of 99 type 1 diabetic patients and 101 controls without diabetes mellitus were included in the study. Sample size of 100 cases and 100 controls was based on the assumption that cases would be 20% less likely to have been breastfed (40% cases vs 60% controls) or 20% more likely to have been bottle fed with (80% cases vs 60%), a power of 0.8 and alpha of 0.05.

Cases were defined as children diagnosed with type 1 diabetes under age of 14 years old attending pediatric endocrine clinic at NGH, Riyadh. Equal number of non diabetic control child were selected from children attending Health Care Specialized Center (HCSC) primary health care in NGH in Riyadh, systematically chosen by a random number from the daily appointment list. Adult and patients with type 2 DM were excluded.

Data on relevant exposures were asked from mothers by means of extensive interview done by a research coordinator to fill the questionnaire that is identical for case and control subjects.

The questionnaire was adopted from a similar study done in Germany<sup>18</sup> after taking permission from its primary author and was customized according to our culture and was validated by an expertise then it was translated to Arabic and back to English.

Basic information was collected on sex, age and diet. History of diet was taken through questions on the duration of overall breastfeeding and age at first introduction of breast milk substitutes, age at introduction of solid foods to infant feeding, type of breast milk substitute fed during the first year of life and current level of customary fresh cow's milk intake.

Mother's age at birth of the index child, child weight at birth, birth order, and the child's medical history were recorded. Further questions addressed family size. Genetic predisposition to of T1DM was covered through family history of T1DM.

Permission was procured from the king Abdullah international medical center in Riyadh, and a verbal consent of each participant `parents obtained before filling the questionnaire.

Statistical package for social sciences (SPSS) software version 19.0 was used for data entry

and analysis. Descriptive statistics (e.g., number and percentage) were calculated for each and every variable wherever applicable. To see the significant difference between the two groups for the continuous variable, we had applied Student's T-test (unpaired). Chi-Square tests ( $\chi^2$ ) were employed to test for the association between two categorical variables we had applied Chi-square test. P values of 0.05 or less were considered statistically significant.

## Results

A total of 200 mothers of children below 14 year old including 101 controls and 99 type 1 diabetic patients were interviewed as shown in table 1. It illustrates that 51.2 % of controls were males and 48.8 % were females and 50 % of cases are female and 50 % were males. Subjects were divided into 4 groups depending on mothers' education level, namely illiterate, primary/intermediate, secondary and university. No significant difference was observed in the distribution of patient or control subjects at any level. It was noticed that children born to mothers with high education level were at high risk of developing diabetes as compared to those who were born to illiterates, although it was not statistically significant ( $p= 0.17$ ). Table 1 also illustrates that no significant difference exists between the patients and controls with regard to the history of bronchial asthma and eczema. Similarly, working status of the parents also did not differ significantly between the two groups.



Figure 1 shows that there is a positive association between birth weight and development of the T1DM. The patients group

tend to be heavier than those with no diabetes are and the difference was statistically significant ( $p=0.009$ ).

*Table 1: Baseline demographic characteristics of patients and control subjects*

Item	No	Cases		Controls		P-value
		No	%	No	%	
Age, years (Means, SD)	<b>200</b>	8.86±2.80		6.33±2.22		<b>0.000</b>
Gender						
Female	<b>114</b>	57	<b>50.0</b>	57	<b>50.0</b>	<b>0.871</b>
Male	<b>86</b>	42	<b>48.8</b>	44	<b>51.2</b>	
Child weight in gm (Means, SD)	<b>200</b>	3140.9±561.3		2929.7±564.4		<b>0.009</b>
Mother with DM						
Yes	<b>16</b>	4	<b>25.0</b>	12	<b>75.0</b>	<b>0.039</b>
No	<b>183</b>	95	<b>51.9</b>	88	<b>48.1</b>	
Mother with GDM						
Yes	<b>17</b>	7	<b>41.2</b>	10	<b>58.8</b>	<b>0.434</b>
No	<b>180</b>	92	<b>51.1</b>	88	<b>48.9</b>	
Mother age at child birth						
≤ 20	<b>28</b>	15	<b>53.6</b>	13	<b>46.4</b>	<b>0.903</b>
21-25	<b>72</b>	35	<b>48.6</b>	37	<b>51.4</b>	
26-46	<b>99</b>	49	<b>49.5</b>	50	<b>50.5</b>	
Order of birth of child						
First born	<b>46</b>	24	<b>52.2</b>	22	<b>47.8</b>	<b>0.88</b>
Second born	<b>65</b>	31	<b>47.7</b>	34	<b>52.3</b>	
Third or more in birth	<b>87</b>	44	<b>50.6</b>	43	<b>49.4</b>	
Sibling with DM						
Yes	<b>25</b>	19	<b>76.0</b>	6	<b>24.0</b>	<b>0.005</b>
No	<b>175</b>	80	<b>45.7</b>	95	<b>54.3</b>	
Gender of sibling with DM						
Female	<b>14</b>	10	<b>71.4</b>	4	<b>28.6</b>	<b>0.26</b>
Male	<b>10</b>	9	<b>90.0</b>	1	<b>10.0</b>	
No. of sibling ≤ 18 years (mean ±SD)	<b>198</b>	4.32±1.97		3.22±1.60		<b>&lt;0.001</b>
Mother education						
Illiterate	<b>24</b>	8	<b>33.3</b>	16	<b>66.7</b>	<b>0.170</b>
Primary/intermediate	<b>61</b>	30	<b>49.2</b>	31	<b>50.8</b>	
Secondary	<b>61</b>	36	<b>59.0</b>	25	<b>41.0</b>	
University	<b>54</b>	25	<b>46.3</b>	29	<b>53.7</b>	
Mother work status						
Housewife	<b>160</b>	79	<b>49.4</b>	81	<b>50.6</b>	<b>0.710</b>
Employee	<b>36</b>	19	<b>52.8</b>	17	<b>47.2</b>	
Asthmatic						
Yes	<b>28</b>	15	<b>53.6</b>	13	<b>46.4</b>	<b>0.640</b>
No	<b>172</b>	84	<b>48.8</b>	88	<b>51.2</b>	
Eczema						
Yes	<b>7</b>	5	<b>71.4</b>	2	<b>28.6</b>	<b>0.230</b>
No	<b>193</b>	94	<b>48.7</b>	99	<b>51.3</b>	

Table 2: Details of breast-feeding and consumption of food among the patients and control subjects

Item	No	Case		Control		P-value
		No	%	No	%	
<b>Breastfeeding given</b>						
No	<b>34</b>	14	<b>41.2</b>	20	<b>58.8</b>	<b>0.280</b>
Yes	<b>166</b>	85	<b>51.2</b>	81	<b>48.8</b>	
<b>Bottle feeding given</b>						
No	<b>25</b>	16	<b>64.0</b>	9	<b>36.0</b>	<b>0.115</b>
Yes	<b>91</b>	82	<b>90.1</b>	9	<b>9.9</b>	
<b>Type of breastfeeding</b>						
Breastfeeding only	<b>25</b>	15	<b>60.0</b>	10	<b>40.0</b>	<b>0.231</b>
Bottle feeding only	<b>141</b>	70	<b>49.6</b>	71	<b>50.4</b>	
Mixed feeding	<b>32</b>	12	<b>37.5</b>	20	<b>62.5</b>	
<b>Duration of breastfeeding</b>						
≤ 4 months	<b>69</b>	34	<b>49.3</b>	35	<b>50.7</b>	<b>0.742</b>
5-6 months	<b>33</b>	16	<b>48.5</b>	17	<b>51.5</b>	
> 6 months	<b>64</b>	35	<b>54.7</b>	29	<b>45.3</b>	
<b>Age at introduction of solid food</b>						
3-4 months	<b>37</b>	19	<b>51.4</b>	18	<b>48.6</b>	<b>0.480</b>
5-6 months	<b>100</b>	65	<b>65.0</b>	35	<b>35.0</b>	
≥ 7 months	<b>20</b>	7	<b>35.0</b>	13	<b>65.0</b>	
<b>Current cow milk consumption</b>						
Non	<b>22</b>	8	<b>36.4</b>	14	<b>63.6</b>	<b>&lt;0.001</b>
< 200 ml	<b>66</b>	27	<b>40.9</b>	39	<b>59.1</b>	
≥ 200 ml	<b>111</b>	64	<b>57.7</b>	47	<b>42.3</b>	

There was no difference between the 2 groups with regard to intake or duration of breastfeeding ( $p= 0.28$ ) as shown in Table 2. It also shows that there was no significant difference between the 2 groups with regard to the age of initiation of formula's milk and weaning. Interestingly we have noticed that with

postponing formula's milk introduction to children, less cases of type 1 DM was diagnosed. Only 25 % of the cases developed if formula's milk postpone until after age of 7 months as compared to 57 % if it introduced before 4 months, however it was not statistically significant ( $p= 0.48$ ).




Fig. 1: Association of T1DM with birth-weight among cases and controls.




Fig. 2: Cases and controls current daily cow's milk consumption.




Fig. 3: Cases and controls with siblings with T1DM

Similar finding with solid food introduction with 35 % of cases of T1DM was diagnosed if it was after age of 7 months as compared to 52 % if started earlier but it was not statistically significant ( $p=0.14$ ). The age at which the subjects were fed solid food for the first time did not differ significantly between the 2 groups ( $p=0.14$ ). The study results showed in Fig. 2 that the current daily consumption of cow's milk in subjects with diabetes were higher significantly in patients group ( $p=0.04$ ).

Interestingly, it was found that children with T1DM were less likely to have a mother with diabetes than controls ( $p=0.039$ ) as seen in Table 1. Also patients with T1DM were more likely to have a sibling with T1DM as over 75 % of patient with T1DM have sibling with DM as compared to 24 % in normal controls ( $p=0.005$ ) as shown in figure 3.

## Discussion

This case control study in children attending NGHHA showed no statistically significant association between breastfeeding, bottle feeding, age at introduction of solid food, or mother age at childbirth, was found with T1DM in children. Factors associated with type 1 diabetes in children were found to be family history of T1DM in siblings, larger birth weight, and high current cow's milk.

Infant's diet has been hypothesized to be involved in the initiation of the T1DM autoimmune process by impairing the

maturation of the gut-associated immune system and or by providing antigens cross-reactive to islet cell antigens (molecular mimicry).<sup>19</sup> The results of this study on infant diet didn't support any protective role of breastfeeding or a late introduction of breast milk substitute as it didn't find any association between duration of breastfeeding between the 2 groups. The diabetogenic effect of animal milk is also a matter of debate. Earlier retrospective studies did not consistently find evidence for a protective effect of breastfeeding and late exposure to formulas or cow's milk.<sup>21-24</sup>

Recent prospective investigations, which were not subject to recall bias, also produced conflicting evidence on the association between infant diet and the risk of beta cell immunity. Some studies did not find an association between diabetes autoimmunity and breastfeeding duration or early formula's milk introduction, whereas other studies did.<sup>25-29</sup> Recently, a randomized double-blinded dietary intervention pilot trial in newborns genetically at increased risk for T1DM provided first evidence that casein hydrolysate formulas may protect against the development of islet cell autoimmunity.<sup>33</sup>

Recent prospective studies found association between increased beta cell autoimmunity and early or late introduction of cereals /gluten or early introduction of fruits and roots into infant diet.<sup>26-28</sup> Both early and late first exposure to any solid food predicted development of T1DM

was shown in one cohort study done in Colorado.<sup>34</sup> The data in this issue is conflicting. The present study did not find any association between the early introduction of solid food and the development of the type 1 diabetes.

Cow's milk influences the composition of the gut flora,<sup>35</sup> In addition cow's milk contains bovine IgG and IgA which could further modify the flora and modulate the immune response to the flora.<sup>36</sup> In the current study, it was noticed that there is significant increase in current daily cow's milk intake inpatient with diabetes. The result of the present study was contrary to the result of one study done in Germany<sup>18</sup> where it showed an inverse relationship in this issue.

A positive family history of T1DM has consistently been reported to raise type 1 diabetes risk among relatives.<sup>20, 37-43</sup> It is known that sibling of a diabetic probands have a higher risk of T1DM than unrelated individuals in the general population at about 3-10%.<sup>43-46</sup> The current study evaluated risk of T1DM in family member and found that T1DM in siblings determines a 3.7 times higher risk of the disease. This is similar to what has been found in study done in Lancashire and Cambria, UK.<sup>36</sup>

The study results showed more cases of maternal T2DM among controls than among cases of T1DM, which support the evidence that there is no association between T1DM and T2DM. This is consistent with studies done in UK<sup>36</sup> and Italy<sup>47</sup> which showed that family

history of type 2 doesn't influence the risk of T1DM in first degree relative.

Epidemiological evidence on the association between level of education of parents and T1DM risk in children is conflicting. In the present study, parents educational background didn't vary between patients and control although there was an interesting finding that mothers with higher education level were having more chance to have diabetic child but it wasn't statistically significant. If we increase the sample size, it may become significant.

Crowded household which was reflected by number of sibling has been observed to be associated with reduced risk for T1DM in case control and recent cohort studies,<sup>20,48-50</sup> Our study did not show any association between the number of sibling and the development of T1DM. The study results did not support the hygiene hypothesis which suggesting that more siblings could lead to earlier and more antigens exposure in life. The hygiene hypothesis suggests that improved hygiene and living conditions have decreased the frequency of childhood infections, leading to a modulation of the developing immune system and increasing risk for autoimmune such as T1DM.<sup>51,52</sup>

There is a considerable body of evidence that higher maternal age at childbirth is associated with a higher risk of T1DM.<sup>39,40,49,53-57</sup> The present study results did not show any association in this regards.

Results in the relation between birth weight and T1DM risk are conflicting. Some studies observed an increased risk in children with high birth weight and lower T1DM risk among children with low birth weight.<sup>57-59</sup> But other studies observed also low birth weight to be associated with increased risk<sup>60-62</sup> while others found no association.<sup>20,55,63</sup> In concordance with the study done in Germany,<sup>18</sup> present study indicated a relationship between higher birth weight and T1DM risk as cases tend to be heavier than control subjects. Data with regard to the order of birth were comparable between the 2 groups in the current study. Other studies have found an increased risk of T1DM among low birth order children.<sup>64-65</sup>

We did not find any significant difference among either sex in developing T1DM which is consistent with one German study<sup>66</sup> which did not find significant difference, whereas the Hawaiian IDDM registry<sup>67</sup> showed a higher incidence among girls. In another study done in India there was a male predominance.<sup>68</sup> Therefore, influence of sex in etiology is not clear.

Regarding the association between atopy and T1DM, one meta-analysis<sup>69</sup> found that there is an inverse relationship between asthma and T1DM. Failure to detect such like association between asthma or eczema and T1DM in the present study may be attributed to the small sample size of our study.



This study has some limitations that should be mentioned. A case control study design generally depends on the collection of retrospective data, thus introducing the possibility of recall bias. The study was conducted in limited area and does not necessary reflect the characteristic of the general population. Finally, there was wide age range (1-14 years). This is because the fact those available cases were limited.

Conclusively, children with T1DM do not differ significantly from their healthy peers in nutritional status. However, child birth-weight, current cow's milk ingestion, and family history of siblings with type 1 diabetes show a significant association with T1DM.

### Acknowledgement

The author would like to express her thanks and appreciation to her academic supervisor Dr. Mazen Ferwana, Associate professor, King Saud bin Abdulaziz University for Health Sciences for the sustained help and expert advice during this study.

### References

1. Lynne L Levitsky, Madhusmita Mirsa. Epidemiology, presentation, and diagnosis of type 1 diabetes mellitus in children and adolescents. <http://www.uptodate.com/contents/epidemiology-presentation-and-diagnosis-of-type-1-diabetes-mellitus-in-children-and-adolescents>
2. Mamoulakis D, Galanakis E, Bicouvarakis S, Paraskakis E, Sbyrakis S. Epidemiology of childhood type I diabetes in Crete, 1990-2001. *Acta Paediatr.* 2003; 92(6):737-9.
3. Karvonen M, Pitkaniemi J, Tuomilehto J. The onset age of type 1 diabetes in Finnish children has become younger. *The Finnish Childhood Diabetes Registry Group. Diabetes Care.* 1999;22(7):1066-70.
4. Scott CR, Smith JM, Craddock MM, Pihoker C. Characteristics of youth-onset noninsulin-dependent diabetes mellitus and insulin-dependent diabetes mellitus at diagnosis. *Pediatrics.* 1997;100(1):84-91.
5. Gale EA, Gillespie KM. Diabetes and gender. *Diabetologia.* 2001;44(1):3-15.
6. Krolewski AS, Warram JH, Rand LI, Kahn CR. Epidemiologic approach to the etiology of type I diabetes mellitus and its complications. *N Engl J Med.* 1987;317(22):1390-8.
7. Patterson CC, Dahlquist GG, Gyürüs E, Green A, Soltész G; EURODIAB Study Group. Incidence trends for childhood type 1 diabetes in Europe during 1989-2003 and predicted new cases 2005-20: a multicentre prospective registration study. *Lancet.* 2009; 373(9680):2027-33
8. Pugliese A. Genetics of type 1 diabetes. *Endocrinol Metab Clin North Am* 2004;33:1-16.
9. Hyttinen V, Kaprio J, Kinnunen L, Koskenvuo M, Tuomilehto J. Genetic liability of type 1 diabetes and the onset age among 22,650 young Finnish twin pairs: a nationwide follow-up study. *Diabetes.* 2003; 52(4):1052-5
10. Nelson Textbook of Pediatrics 19th edition page: Robert M. Kliegman, Bonita F. Stanton, Joseph W. St. Geme III, Nina F. Schor, Richard E. Behrman Chapter 583.2 Type 1 Diabetes Mellitus (Immune Mediated) Ramin Alemzadeh and Omar Ali
11. Savilahti E, Akerblom HK, Tainio V-M, Koskimies S. Children with newly diagnosed insulin dependent diabetes mellitus have increased levels of cow's milk antibodies. *Diabetes Res.* 1988;7(3):137-40.
12. Dahlquist G, Savilahti E, Landin-Olsson M. An increased level of antibodies to beta-lactoglobulin is a risk determinant for early-onset type 1 (insulin-dependent) diabetes mellitus independent of islet cell antibodies and early introduction of cow's milk. *Diabetologia.* 1992;35(10):980-4.
13. Saukkonen T, Savilahti E, Vaarala O, Virtala ET, Tuomilehto J, Akerblom HK. Children with newly diagnosed IDDM have increased levels of antibodies to bovine serum albumin but not to ovalbumin. *Diabetes Care.* 1994;17(9):970-6.
14. Catassi C, Bonucci A, Coppa GV, Carlucci A, Giorgi PL. Intestinal permeability changes during the first month: effect of natural versus artificial feeding. *J Pediatr Gastroenterol Nutr.* 1995 Nov;21(4):383-6.
15. Knip M. Environmental triggers and determinants of b-cell autoimmunity and type 1 diabetes. *Rev Endocr Metab Disord.* 2003;4(3):213-23.

16. Jenista JA, Powell KR, Menegus MA. Epidemiology of neonatal enterovirus infection. *J Pediatr.* 1984;104(5):685-90.
17. Sadeharju K, Knip M, Virtanen SM, Savilahti E, Tauriainen S, Koskela P, et al. Maternal antibodies in breast milk protect the child from enterovirus infections. *Pediatrics.* 2007;119(5):941-6.
18. Rosenbauer J, Herzig P, Giani G. Early infant feeding and risk of type 1 diabetes mellitus—a nationwide population-based case-control study in pre-school children. *Diabetes Metab Res Rev.* 2008;24(3):211-22.
19. Harrison LC, Honeyman MC. Cow's milk and type 1 diabetes the real debate is about mucosal immune function. *Diabetes.* 1999;48(8):1501-7.
20. Wadsworth EJ, Shield JP, Hunt LP, Baum JD. A case-control study environmental factors associated with diabetes in the under 5s. *Diabet Med.* 1997;14(5):390-6.
21. Nigro G, Campea L, DeNovellis A, Orsini M. Breast-feeding and insulin-dependent diabetes mellitus. *Lancet.* 1985;1(8426):467.
22. Siemiatycki J, Colle E, Campbell S, Dewar RA, Belmonte MM. Case-control study of IDDM. *Diabetes Care.* 1989;12(3):209-16
23. Kyvik KO, Green A, Svendsen A, Mortensen K. Breast feeding and the development of type 1 diabetes mellitus. *Diabet Med.* 1992;9(3):233-5.
24. Virtanen SM, Laara E, Hypponen E, Reijonen H, Räsänen L, Aro A, et al. Cow's milk consumption, HLA-DQB1 genotype, and type 1 diabetes: a nested case-control study of siblings of children with diabetes: Childhood diabetes in Finland study group. *Diabetes.* 2000;49(6):912-7.
25. Couper JJ, Steele C, Beresford S, Powell T, McCaul K, Pollard A, et al. Lack of association between duration of breast-feeding or introduction of cow's milk and development of islet autoimmunity. *Diabetes.* 1999;48(11):2145-9.
26. Ziegler AG, Schmid S, Huber D, Hummel M, Bonifacio E. Early infant feeding and risk of developing type 1 diabetes-associated autoantibodies. *JAMA.* 2003;290(13):1721-8.
27. Norris JM, Barriga K, Klingensmith G, Hoffman M, Eisenbarth GS, Erlich HA, et al. Timing of initial cereal exposure in infancy and risk of islet autoimmunity. *JAMA.* 2003;290(13):1713-20.
28. Virtanen SM, Kenward MG, Erkkola M, Kautiainen S, Kronberg-Kippilä C, Hakulinen T, et al. Age at introduction of new foods and advanced beta cell autoimmunity in young children with HLA-conferred to susceptibility to type 1 diabetes. *Diabetologia.* 2006;49(7):1512-21.
29. Kimpimäki T, Erkkola M, Korhonen S, Kupila A, Virtanen SM, Ilonen J, et al. Short-term exclusive breastfeeding predisposes young children with increased genetic risk of Type I diabetes to progressive beta-cell autoimmunity. *Diabetologia.* 2001;44(1):63-9.
30. Wahlberg J, Vaarala O, Ludvigsson J. Dietary risk factors for the emergence of type 1 diabetes-related autoantibodies in 21/2 year-old Swedish children. *Br J Nutr.* 2006;95(3):603-8.
31. Patelarou E, Girvalaki C, Brokalaki H, Patelarou A, Androulaki Z, Vardavas C. Current evidence on the associations of breastfeeding, infant formula, and cow's milk introduction with type 1 diabetes mellitus: a systematic review. *Nutr Rev.* 2012;70(9):509-19
32. Knip M, Virtanen SM, Becker D, Dupré J, Krischer JP, Åkerblom HK, et al. Early feeding and risk of type 1 diabetes: experiences from the Trial to Reduce Insulin-dependent diabetes mellitus in the Genetically at Risk (TRIGR). *Am J Clin Nutr.* 2011;94(6 Suppl):1814S-1820S.
33. Åkerblom HK, Virtanen SM, Ilonen J, Savilahti E, Vaarala O, Reunanen A, et al. Dietary manipulation of beta cell autoimmunity in infants at increased risk of type 1 diabetes: a pilot study. *Diabetologia.* 2005;48(5):829-37.
34. Frederiksen B, Kroehl M, Lamb MM, Seifert J, Barriga K, Eisenbarth GS, et al. Infant Exposures and Development of Type 1 Diabetes Mellitus, The Diabetes Autoimmunity Study in the Young (DAISY). *JAMA Pediatr.* 2013;167(9):808-15.
35. Mackowiak PA. The normal microbial flora. *N Engl J Med.* 1982 8;307(2):83-93..
36. Marshall AL, Chetwynd A, Morris JA, Placzek M, Smith C, Olabi A, et al. Type 1 diabetes mellitus in childhood: a matched case control study in Lancashire and Cumbria, UK. *Diabet Med.* 2004;21(9):1035-40.
37. Gillespie KM, Gale EA, Bingley PJ. High familial risk and genetic susceptibility in early onset childhood diabetes. *Diabetes.* 2002;51(1):210-4.
38. Glatthaar C, Whittall DE, Welborn TA, Gibson MJ, Brooks BH, Ryan MM, et al. Diabetes in Western Australian children: descriptive epidemiology. *Med J Aust.* 1988;148(3):117-23.
39. Dahlquist G, Blom L, Lönnerberg G. The Swedish Childhood Diabetes Study—a multivariate analysis of risk determinants for diabetes in different age groups. *Diabetologia.* 1991; 34(10):757-62.
40. McKinney PA, Parslow R, Gurney KA, Law GR, Bodansky HJ, Williams R. Perinatal and neonatal

- determinants of childhood type 1 diabetes. A case-control study in Yorkshire, U.K. *Diabetes Care*. 1999;22(6):928-32.
41. Sipetić SB1, Vlajinac HD, Kocev NI, Marinković JM, Radmanović SZ, Bjekić MD. The Belgrade childhood diabetes study a multivariate analysis of risk determinants for diabetes. *Eur J Public Health*. 2005;15(2):117-22.
  42. Marshall AL, Chetwynd A, Morris JA, Placzek M, Smith C, Olabi A, et al. Type 1 diabetes mellitus in childhood: a matched case control study in Lancashire and Cumbria, UK. *Diabet Med*. 2004;21(9):1035-40.
  43. Altobelli E1, Chiarelli F, Valenti M, Verrotti A, Blasetti A, Di Orio F. Family history and risk of insulin-dependent diabetes mellitus: a population-based case-control study. *Acta Diabetol*. 1998;35(1):57-60.
  44. Majeed AAS, Hassan MK. Risk factor of type 1 DM among children and adolescents in Basrah. *Oman Med J*. 2011;26:189-195.
  45. Ramachandran A, Snehalatha C, Premila L, Mohan V, Viswanathan M. Familial aggregation in type 1 (insulin-dependent) diabetes mellitus: a study from south india. *Diabetic Medicine* 1990;7(10): 876-879.
  46. Warram JH, Krolewski AS, Gottlieb MS, Kahn CR. Differences in risk of insulin- dependant diabetes in offspring of diabetic mothers and diabetic fathers. *N Engl J Med*. 1984; 311(3):149-52.
  47. Altobelli E, Chirraelli F, Valenti M, Verrotti A, Blasetti A, Di Orio F. Family history and risk of insulin-dependent diabetes mellitus: a population-based case-control study. *Acta Diabetol* 1998; 35(1):57-60.
  48. Verge CF, Howard NJ, Irwig L, Simpson JM, Mackerras D, Silink M. Environmental factors in childhood IDDM. A population-based, case-control study. *Diabetes Care*. 1994; 17(12):1381-9.
  49. Patterson CC, Carson DJ, Hadden DR, Waugh NR, Cole SK. A case-control investigation of perinatal risk factors for childhood IDDM in Northern Ireland and Scotland. *Diabetes Care*. 1994; 17(5):376-81.
  50. Stene LC, Magnus P, Lie RT, Sovik O, Joner G. Maternal and paternal age at delivery, birth order, and risk of childhood onset type 1 diabetes: population based cohort study. *BMJ* 2001; 323-369.
  51. Bach JF. Six questions about the hygiene hypothesis? *Cell Immunol*. 2005;233(2):158-61.
  52. Gale EAM. A missing link in the hygiene hypothesis? *Diabetologia*. 2002;45(4):588-94.
  53. Blom L, Dahlquist G, Nystrom L, Sandstrom A, Wall S. The Swedish childhood diabetes study—social and perinatal determinants for diabetes in childhood. *Diabetologia*. 1989;32(1):7-13.
  54. Soltész G, Jeges S, Dahlquist G. Non-genetic risk determinants for type 1 (insulin-dependent) diabetes mellitus in childhood Hungarian Childhood Diabetes Epidemiology Study Group. *Acta Paediatr*. 1994; 83(7):730-5.
  55. McKinney PA, Okasha M, Parslow RC, Law GR, Gurney KA, Williams R, et al. Early social mixing and childhood Type 1 diabetes mellitus: a case- control study in Yorkshire, UK. *Diabet Med*. 2000; 17(3): 236-42.
  56. Patterson CC, Dahlquist G, Soltesz G. Maternal age and risk of type 1 diabetes in children. Relative risks by maternal age are biased. *BMJ*. 2001;322(7300):1489-90
  57. Kostraba JN, Cruickshanks KJ, Lawler-Heavner J, Jobim LF, Rewers MJ, Gay EC, et al. Early exposure to cow's milk and solid foods in infancy, genetic predisposition, and risk of IDDM. *Diabetes*. 1993;42(2):288-95.
  58. Dahlquist GG, Patterson C, Soltesz G. Perinatal risk factors for childhood type 1 diabetes in Europe. The EURODIAB Substudy 2 Study Group. *Diabetes Care*. 1999;22(10):1698-702.
  59. Stene LC, Magnus P, Lie RT, Sjøvik O, Joner G; Norwegian childhood Diabetes Study Group. Birth weight and childhood onset type 1 diabetes: population based cohort study. *BMJ*. 200;322(7291):889-92.
  60. Tai TY, Wang CY, Lin LL, Lee LT, Tsai ST, Chen CJ. A case control study on risk factors for Type 1 diabetes in Taipei City. *Diabetes Res Clin Pract*. 1998;42(3):197-203.
  61. Stene LC, Thorsby PM, Berg JP, Rønningen KS, Undlien DE, Joner G, et al. The relation between size at birth and risk of type 1 diabetes is not influenced by adjustment for the insulin gene(-23HphI) polymorphism or HLA-DQ genotype. *Diabetologia*. 2006;49(9):2068-73
  62. Wei JN, Li HY, Chang CH, Sung FC, Li CY, Lin CC, et al. Birth weight and type 1 diabetes among schoolchildren in Taiwan—A population-based case controlled study. *Diabetes Res Clin Pract*. 2006;74(3):309-15.
  63. Jones ME, Swerdlow AJ, Gill LE, Goldacre MJ. Pre-natal and early life risk factors for childhood onset diabetes mellitus: a record linkage study. *Int J Epidemiol*. 1998;27(3):444-9.
  64. Metcalfe MA, Baum JD. Family characteristics and insulin dependent diabetes. *Arch Dis Child*. 1992;67(6):731-6.
  65. Ramachandran A1, Snehalatha C, Joseph TA, Vijay V, Viswanathan M. Delayed onset of diabetes in children

- of low economic stratum- A study from South India. *Diabetes Res Clin Pract.* 1994;22(2-3):171-4.
66. Nek A, Hub R, Kehren M, Ranke MB. Incidence of IDDM in German children aged 0-14 years-a 6 year population study. *Diabetes Care.* 1997;20(4):530-3.
67. Schoenle EJ, Lang-Muritano M, Gschwend S, Laimbacher J, Mullis PE, Torresani T, et al. Epidemiology of type I diabetes mellitus in Switzerland: Steep rise in incidence under 5 years old children in past decades. *Diabetologia.* 2001;44(3):286-9.
68. Manash P, Ariachery C, Madan L. Demographic, breastfeeding , and nutritional trends among children with type 1 diabetes mellitus. *Indian J Endocrinol Metab.* 2011;15(1):38-42.
69. Chris R, Mike D, Dennes J and Chris C. A meta-analysis of the association between childhood type 1 diabetes and atopic disease. *Diabetes Care.* 2003;26(9):2568-74.

**ORIGINAL ARTICLE**

## Prevalence and Antibiotic Resistance of Cronobacter Spp. Associated with Powdered Infant Milk Formulas and Dried Milks in Saudi Arabia

Khaled M. Aljarallah\*<sup>1</sup>

\* For correspondence [k.aljarallah@mu.edu.sa](mailto:k.aljarallah@mu.edu.sa)

<sup>1</sup>Department of Medical Laboratory Sciences, College of Applied Medical Sciences, Majmaah University, Almajmaah-11952, Kingdom of Saudi Arabia

Received on: 13<sup>th</sup> October 2015, Accepted on: 27<sup>th</sup> October 2015

### Abstract

**Introduction:** This study examined the presence of the pathogen Cronobacter spp. in powdered infant milk formulas (PIMF), and dried whole and skim milk samples distributed in Saudi Arabia. **Methodology:** Culture-based method, biochemical conformation test and real-time PCR were used for detection. **Results:** Cronobacter spp. was detected in 10% of PIMF and 20% of dried whole milk, but not in dried skim milk samples. Real-time PCR showed that PIMF samples generally contained higher numbers of Cronobacter spp. than dried whole milk samples. Isolates showed resistance to 9 of 10 antibiotics, being more frequently resistant to penicillin and tetracycline Both single and multidrug resistance was observed. **Conclusions:** This study reports the association of antibiotic-resistant Cronobacter spp. with PIMF and dried whole milk products distributed in Saudi Arabia.

**Key Words:** Cronobacter, antibiotic resistance, infant milk formulas, real-time PCR

### المخلص

**مقدمة:** تناولت هذه الدراسة مدى تواجد بكتيريا الكرونوباكتر (Cronobacter spp.) الممرضة في عينات المركبات الحليبية للأطفال الرضع إضافة لعينات حليب مجفف كامل ومنزوع الدسم والتي يتم تداولها في المملكة العربية السعودية. تم الكشف عن تواجد هذا الميكروب الممرض في العينات باستخدام عدة طرق شملت الاستزراع والفحص الكيموحيوي التأكيدي إضافة للفحص الجزيئي باستخدام تفاعل إنزيم البلمرة التسلسلي الأني (Real time PCR). تم الكشف عن تواجد بكتيريا الكرونوباكتر في 10% من عينات المركبات الحليبية للأطفال الرضع و 20% من عينات الحليب المجفف كامل الدسم بينما خلت منه عينات الحليب المجفف منزوع الدسم. بينت نتائج الفحص باستخدام إنزيم البلمرة التسلسلي الأني (Real time PCR) أن عينات المركبات الحليبية للأطفال الرضع تحتوي في الغالب أعدادا أكبر من هذا الجنس البكتيري مقارنة بعينات الحليب المجفف كامل الدسم. إختبارات المضادات الحيوية أظهرت أن البكتيريا المعزولة في هذه الدراسة كانت مقاومة لتسع مضادات حيوية من أصل عشرة تم استخدامها , وكانت المقاومة الأكثر تكرارا ووضوحا ضد البنيسلين والتتراسايكلين . المقاومة الأحادية والمتعددة كلاهما تم ملاحظته . هذه الدراسة أثبتت تواجد بكتيريا الكرونوباكتر المقاومة للمضادات الحيوية في مركبات حليبية للأطفال الرضع وكذلك في منتجات حليب مجفف كامل الدسم تسوق في المملكة العربية السعودية.

**Introduction:** Cronobacter spp. (formerly known as Enterobacter sakazakii) is an opportunistic pathogen that has emerged recently in association with sporadic disease cases and outbreaks involving debilitated neonates fed dried infant formulas.<sup>1-3</sup> Infections with Cronobacter spp. may lead to meningitis, necrotising enterocolitis, and bacteraemia that could be associated with a high mortality rate of 40–80%<sup>4</sup> Brain damage or neurological

disorders, such as hydrocephalus and quadriplegia, may also manifest in surviving patients<sup>5</sup>. Both infants and elderly individuals were found to be infected by Cronobacter spp.<sup>6</sup> While the pathogen was recovered from various sources, including soil, water, vegetables, and foods, it was most frequently detected in dried milks and powdered infant milk formulas (PIMF).<sup>7-11</sup> Cronobacter spp. has thus been classified as a category A organism that is



clearly associated with causing disease via PIMF.<sup>3</sup> Previous studies have reported the association of *Cronobacter* spp. with dried milks and PIMF in different Western countries. Other reports also demonstrated its presence in PIMF and dried milk products distributed in the Middle and Far Eastern regions, including Jordan,<sup>12-13</sup> Egypt<sup>10,14</sup> and China.<sup>15</sup> A previous study also showed the contamination of the Saudi Arabian artisanal fermented drink "Sobia," made from wheat and malt flours, with *Cronobacter* spp.<sup>16</sup> However, no further information is available on the association of *Cronobacter* spp. with PIMF and powdered milks in Saudi Arabia. The present study was therefore designed to address this aspect, and to characterise the antibiotic resistance of *Cronobacter* spp. isolates associated with these products.

## Methodology

Detection of *Cronobacter* spp. in powdered infant milk formulas and dried milk samples: A total of 60 samples of powdered infant milk formulas (PIMF) (n=20), dried whole milk (n=20), and dried skim milk (n=20) were randomly collected from local markets in Riyadh, Saudi Arabia. These samples were examined for the presence of *Cronobacter* spp. using the FDA method<sup>8</sup>, as modified by El-Sharoud et al.<sup>10,14,11</sup>. Samples were pre-enriched by mixing a 25 g sample with 225 mL buffered peptone water (BPW) broth (Oxoid, Basingstoke, UK), followed by incubation at

37°C for 24 h. Aliquots of 10 mL of the pre-enriched samples were then inoculated into 90 mL of the EE broth (Oxoid), followed by incubation at 37°C for 24 h. A 3 mm loopful (10 µL) of the enterobacteriaceae enrichment (EE) broth culture was finally streaked onto the brilliance *Enterobacter sakazakii* agar (DFI agar) (Oxoid) followed by incubation at 37°C for 24 h. Suspected blue-green colonies were then picked up and examined by Gram-staining and for the formation of yellow pigmented colonies on tryptone soy agar (TSA) (Oxoid) at 25°C for 48–72 h. Potential isolates producing a negative Gram-reaction and yellow pigment on TSA were further examined using the Rapid ID 32 E miniaturised kit (bioMerieux, Marcy l'Etoile, France).

*Real-time PCR confirmation of Cronobacter spp.:* Real-time PCR was used to confirm *Cronobacter* spp. isolates recovered from the PIMF and dried milk samples. The real-time PCR protocol<sup>17</sup> of was applied as follows. Cells were pelleted from a 24 h culture of each isolate by centrifugation at 5000 rpm for 10 min. The cell pellet was re-suspended in PrepMan Ultra (Applied Biosystems, Foster City, CA, USA) and heated at 100°C in a thermal block incubator for 15 min, followed by cooling for 2 min at room temperature, and centrifugation at 5000 rpm for 2 min. Resultant supernatants containing DNA were collected and used for real-time PCR reactions. A PCR reaction mixture of a total volume of 50 µL was

formulated using a 5  $\mu$ L DNA sample template, 25  $\mu$ L of TaqMan Universal PCR Master Mix (Applied Biosystems), 5  $\mu$ L of forward primer, 5  $\mu$ L of reverse primer, 5  $\mu$ L of TaqMan probe, and 5  $\mu$ L of water. The primers and TaqMan probe were designed to target a DNA sequence of 78 bp within the macromolecular synthesis (mms) operon<sup>17</sup> for 2 min, followed by 95 °C for 10 min, and 50 cycles of 95 °C for 15 s and 60 °C for 60 s. PCR reactions were conducted using a StepOne real-time PCR system (Applied Biosystems).

*Assessment of the antibiotic resistance of Cronobacter spp.:* Cronobacter spp. isolates were examined for their resistance to 10 antibiotics, including ampicillin (10  $\mu$ g), penicillin G (100 unit), gentamicin (10  $\mu$ g), tetracycline (30  $\mu$ g), ciprofloxacin (5  $\mu$ g), kanamycin (30  $\mu$ g), streptomycin (10  $\mu$ g), chloramphenicol (30  $\mu$ g), nalidixic acid (30  $\mu$ g), and cefoxitin (30  $\mu$ g). Antibiotic resistance was tested using the Kirby-Bauer disc-diffusion method. Briefly, a standardised inoculum of a 24 h culture of each isolate was spread onto Muller-Hinton agar (Oxoid), followed by dispensing antibiotic-impregnated discs. After incubation at 37° C for 24 h, the diameter of the inhibition zones around each antibiotic disc was measured and interpreted as resistance, intermediate, or sensitive, according to the criteria of the Clinical and Laboratory Standards Institute.<sup>18</sup>

## Results

Presence of Cronobacter spp. in powdered infant milk formulas and dried milk products. The prevalence of Cronobacter spp. in powdered infant milk formulas (PIMF) and dried whole and skim milk samples collected in March 2013 from the local markets in Saudi Arabia was examined.

*Table 1: Detection of Cronobacter spp. in powdered infant milk formulas and dried milk samples*

Product	No. of samples	No. of presumptive Cronobacter spp. positive isolates	No. of bio-chemically confirmed isolates (%)*	No. of real-time PCR confirmed isolates (%)*	Minimum Ct	Maximum Ct	No. of positive samples (%)
Powdered infant milk formulas	20	5	2 (40)	2 (40)	15.8	20	2 (10)
Dried whole milk	20	10	4 (40)	4 (40)	19.8	25	4 (20)
Dried skim milk	20	0	0	0	-		0
Total	60	15	6 (40)	6 (40)			6 (10)

A total of 15 potential isolates of the organism producing typical colonies on the DFI agar, a negative Gram reaction, and a yellow pigment on the TSA were recovered from these samples (Table 1).

However, only 40% of these isolates could be confirmed by biochemical testing using the miniaturised rapid ID 32E kit and a real-time PCR assay targeting the mms operon (Table 1 and Figure 1).




Figure 1: Real-time PCR analysis of *Cronobacter* spp. Isolates.

Figure 1 shows real-time PCR curves resulting from the analysis of the *Cronobacter* spp. isolates recovered from the PIMF and whole milk powder samples. Ct number is inversely related to the amount of amplicon in the reaction (target) whereas the lower the Ct, the greater the amount of amplicon<sup>9</sup>. This was reflected here where the number of isolates in the powdered infant milk formulas and dried whole milk were 2 and 4 which correspond to the Ct values 17.9 and 22.4 respectively.




Figure 2: Response of *Cronobacter* spp. Isolates. Isolates showed variable rates of resistance (■), intermediate response (▒), and sensitivity (□) to the examined antibiotics.

Antibiotic resistance of *Cronobacter* spp. Isolates: Among the examined products, 10% and 20% of PIMF and dried whole milk samples were found to be contaminated with *Cronobacter* spp., respectively (Table 1). Dried skim milk samples were found free of the organism.

The susceptibility of *Cronobacter* spp. isolates recovered from PIMF and dried whole milk samples to the 10 antibiotics was examined. With the exception of gentamycin, *Cronobacter* spp. isolates showed variable resistance to the examined antibiotics (Figure 2).

## Discussion

The highest resistance was found against penicillin, followed by tetracycline, with a resistance rate of 85% and 70% of the isolates, respectively. Resistance to other antibiotics ranged from 10% to 60% of the isolates. These results were consistent with those of Farmer et al.<sup>19</sup>, who reported resistance to penicillin in all examined *Cronobacter* spp. cultures isolated from clinical samples.

Kilonzo-Nthenge et al.<sup>20</sup> also showed that the highest antibiotic resistance rates in *Cronobacter* spp. isolates recovered from domestic kitchens were found against penicillin followed by tetracycline. The authors have also found the isolates to be susceptible to gentamycin. However, they reported a higher resistance to ciprofloxacin of 57% of the isolates compared to the present study (15%), reflecting the effect of

the isolate source on its antibiotic resistance pattern.

*Cronobacter* spp. isolates recovered from PIMF and dried whole milk samples examined in this study showed both single antibiotic resistance and multidrug resistance to more than two antibiotics. Multidrug resistance was observed against penicillin, tetracycline, and streptomycin. Previous studies demonstrated multidrug resistance to various combinations of antibiotics in *Cronobacter* spp. isolates cultured from an adult patient with a wound infection and from domestic kitchens.<sup>20-21</sup>

Despite the assumption that *Cronobacter* spp. infections can be eradicated using antibiotics, the present study confirms the results of other previous reports showing that the pathogen may be able to resist several effective antibiotics. This raises concern regarding the use of antibiotics to treat these infections that could possibly lead to serious illnesses, including meningitis. Drug resistance in foodborne bacteria has been frequently linked to the abuse of antibiotics in feeding animals.<sup>22</sup> In addition to complicating medical treatments, antibiotic-resistant foodborne pathogens could serve as a reservoir spreading genetic elements of antibiotic resistance to other bacteria in the human gut.<sup>23</sup>

These results indicated the importance of employing biochemical and molecular identification testing for reliable detection of *Cronobacter* spp. This is consistent with

previous reports demonstrating the value of incorporating the rapid ID 32E system and real-time PCR analysis in the detection protocols of *Cronobacter* spp. in PIMF and other related products.<sup>24-25</sup>

## Conclusion

In conclusion, the present study demonstrates the association of the emerging pathogen *Cronobacter* spp. with PIMF and whole dried milk in Saudi Arabia. It also reports variability in the prevalence rates of the pathogen in these products compared to the rates in other regions. The study highlights the ability of *Cronobacter* spp. isolates to develop both single and multidrug resistance, which raises concern regarding the effectiveness of available antibiotics to combat serious illnesses associated with *Cronobacter* spp. infection. This requires further research in order to develop more effective antibiotics and/or alternative therapeutic strategies.

## References

1. Gurtler J.B. & Beuchat L.R. Performance of media for recovering stressed cells of *Enterobacter sakazakii* as determined using spiral plating and ecometric techniques," *Applied and Environmental Microbiology*, 2005:71, 7661–7669.
2. Mullane N.R., Iversen C., Healy B., Walsh C., Whyte P., Wall P.G., Quinn T., Fanning S. *Enterobacter sakazakii* an emerging bacterial pathogen with implications for infant health, *Minerva Pediatrica*, 2007:59, 137–148.
3. Forsythe S.J., Dickins B., Jolley K.A. *Cronobacter* spp. spp., the emergent bacterial pathogen *Enterobacter sakazakii* comes of age; MLST and whole genome sequence analysis," *BMC Genomics*, 2014: doi:10.1186/1471-2164-15-1121.

4. Farber J.M. Enterobacter sakazakii – new foods for thought?, *Lancet*, 2004;363, 5–6.
5. Biering G., Karlsson S., Clark N.C., Jónsdóttir K.E., Lúdvígsson P., Steingrímsson O. Three cases of neonatal meningitis caused by Enterobacter sakazakii in powdered milk, *Journal of Clinical Microbiology*, 1989;27, 2054–2056.
6. Tall B.D., Chen Y., Yan Q., Gopinath G.R., Grim C.J., Jarvis K.G., Fanning S., Lampel K.A. Cronobacter spp. spp.: An emergent pathogen causing meningitis to neonates through their feeds, *Science Progress*, 2014;97, 154–172.
7. Iversen C., Lane M., Forsythe S.J. The growth profile, thermotolerance and biofilm formation of Enterobacter sakazakii grown in infant formula milk, *Letters in Applied Microbiology*, 2004;38, 378–382.
8. Nazarowec-White M. & Farber J.M. Prevalence, survival, and growth of Enterobacter sakazakii in infant formula, *Journal of Food Protection*, 1997;60, 226–230.
9. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method. *Nat Protoc.*2008;3:1101–1108. doi: 10.1038/nprot.2008.73.
10. El-Sharoud W.M., El-Din M.Z., Ziada D.M., Ahmed S.F., Klena J.D. Surveillance and genotyping of Enterobacter sakazakii suggest its potential transmission from milk powder into imitation recombined soft cheese, *Journal of Applied Microbiology*, 2008;105, 559–566.
11. El-Sharoud W.M., Darwish M. S., Batt C.A. A real-time PCR-based microfluidics platform for the detection of Cronobacter spp. spp. sakazakii in reconstituted milks, *International Dairy Journal*, 2013;33, 67–74.
12. Shaker R., Osaili T., Al-Omary W., Jaradat Z., Al-Zuby M. Isolation of Enterobacter sakazakii and other Enterobacter sp. from food and food production environments, *Food Control*, 2007;18, 1241–1245.
13. Jaradat Z.W., Ababneh Q.O., Saadoun I.M., Samara N.A., Rashdan A.M. Isolation of Cronobacter spp. spp. (formerly Enterobacter sakazakii) from infant food, herbs and environmental samples and the subsequent identification and confirmation of the isolates using biochemical, chromogenic assays, PCR and 16S rRNA sequencing," *BMC Microbiology*, 2009;doi:10.1186/1471-2180-9-225.
14. El-Sharoud W.M., O'Brien S., Negredo C., Iversen C., Fanning S., Healy B. Characterization of Cronobacter spp. spp. recovered from dried milk and related products, *BMC Microbiology*, 2009;9, doi 10.1186/1471-2180-9-24.
15. Cui J., Du X., Liu H., Hu G., Lv G., Xu B., Yang X., Li W., Cui Z. The genotypic characterization of Cronobacter spp. spp. isolated in China," *PLoS ONE*, 2014;9, doi:10.1371.
16. Gassem M.A.A. A microbiological study of Sobia: A fermented beverage in the Western province of Saudi Arabia, *World Journal of Microbiology and Biotechnology*, 2002;18, 173–177.
17. Seo K.H. & Brackett R.E. (2005): Rapid, specific detection of Enterobacter sakazakii in infant formula using a real-time PCR assay, *Journal of Food Protection*, 68, 59–63.
18. Clinical and Laboratory Standards Institute (CLSI) (2011): Performance standards for antimicrobial susceptibility testing, Wayne, PA, USA, CLSI, M 100–S21.
19. Farmer J.J., Asbury M.A., Hickman F.W., Brenner D.J. & The Enterobacteriaceae Study Group. Enterobacter sakazakii, new species of Enterobacteriaceae isolated from clinical samples, *International Journal of Systematic Bacteriology*, 1980;30, 569–584.
20. Kilonzo-Nthenge A., Rotich E., Godwin S., Nahashon S., Chen F. (2012): Prevalence and antimicrobial resistance of Cronobacter spp. spp. sakazakii isolated from domestic kitchens in Middle Tennessee, United States, *Journal of Food Protection*, 75, 1512–1517.
21. Dennison S.K. & Morris J. (2002): Multiresistant Enterobacter sakazakii wound infection in an adult, *Infection and Medicine*, 2001;1,533–535.
22. McEwen S.A. & Fedorka-Cray P.J. Antimicrobial use and resistance in animals," *Clinical and Infectious Diseases*, 2002;34,S93–S106.
23. El-Sharoud W.M., Yassin M.A., Ahmed S.F. (2015): Molecular characterisation and stress tolerance of Escherichia coli isolated from dairy and dried milk-related products, *International Journal of Food Science and Technology*, 50, 136–142.
24. Lampel K.A. & Chen Y. (2009): Method for the isolation and detection of Enterobacter sakazakii (Cronobacter spp. spp.) from powdered infant formula," *International Journal of Food Microbiology*, 136, 179–184.
25. Chen Y., Song K.Y., Brown E.W., Lampel K.A. (2010): Development of an improved protocol for the isolation and detection of Enterobacter sakazakii (Cronobacter spp. spp.) from powdered infant formula," *Journal of Food Protection*, 73, 1016–1022.



**ORIGINAL ARTICLE**

## Fast Food Consumption among University Students, Saudi Arabia A Cross Sectional Study

Fahad AlFahied<sup>\*1</sup>, Waqas Sami<sup>2</sup>, Yasser AlTowyan<sup>3</sup>, Turki AlJohani<sup>3</sup>, Abdulaziz AlHaddad<sup>3</sup>,  
Abdullah AlKhtami<sup>3</sup>

\* For correspondence [f.alfahed@mu.edu.sa](mailto:f.alfahed@mu.edu.sa)

<sup>1</sup>Assistant Professor, Family Medicine, College of Medicine, Majmaah University, Al Majmaah, KSA

<sup>2</sup>Lecturer, Biostatistics, College of Medicine, Majmaah University, Al Majmaah, KSA

<sup>3</sup>Medical students, College of Medicine, Majmaah University, Al Majmaah,

Received on: 27<sup>th</sup> May 2015, Accepted on: 6<sup>th</sup> September 2015

### Abstract

**Introduction:** Fast food consumption (FFC) is one of the most common eating habits nowadays with escalated prevalence in the recent decades. The study aimed to find the prevalence of FFC among the University students, to identify the factors that contributes to the FFC among them, and to test student's level of awareness regarding the health issues occur due to the consumption of fast food. **Methodology:** A total of 505 students (325) males and (180) females aged between 18-26 years were randomly chosen from Colleges of Majmaah University. **Results.** 41.6% of the participants choose fast food as their usual meal. Male students consumed fast food as a usual meal more and female students were attracted more by taste and advertisements. Consuming fast food and staying away from the family was significantly more in male students as compared to female students. **Conclusions:** The prevalence of fast food consumption among Majmaah University students is very high.

**Key Words:** Fast food consumption, University students, World health organization, electronic media.

### المخلص

**مقدمة:** تناول الوجبات السريعة هي واحدة من العادات الغذائية الأكثر شيوعاً في الوقت الحاضر مع تصاعد ملحوظ في العقود الأخيرة. وتعد الوجبات السريعة أحد عوامل الخطر للعديد من الأمراض، وقد لوحظ ان الخطر يزداد مع الزيادة في استهلاك الوجبات السريعة. الأهداف: تهدف هذه الدراسة لمعرفة نسبة انتشار تناول الوجبات السريعة بين طلاب جامعة المجمعة، كذلك تحديد العوامل التي تساهم في استهلاكها بينهم، واختبار مستوى وعي الطالب بشأن الاضرار الصحية التي تحدث نتيجة لاستهلاك الوجبات السريعة. **الطريقة:** هذه دراسة مقطعية من ديسمبر 2013 -مايو 2014. تشمل عينة الدراسة الطلاب الذين يدرسون في جامعة المجمعة من الجنسين. شملت الدراسة ما مجموعه 505 طالبا (325) من الذكور و (180) من الإناث الذين تتراوح أعمارهم بين 18-26 عاما وقد تم اختيارهم عشوائيا من كليات جامعة المجمعة **النتائج:** مائتان وعشرة (41.6%) من المشاركين (159 ذكور، 51 إناث) يعتبر الوجبات السريعة هي الوجبة اليومية المعتادة. وقد لوحظ ارتفاع هذه النسبة من قبل الطلاب بالمقارنة مع الطالبات ( $P < 0.001$ ). كذلك بينت الدراسة ان الاعلانات تجذب الطالبات للوجبات السريعة أكثر من الطلاب ( $P < 0.001$ ). **الاستنتاج:** ارتفاع معدل تناول الوجبات السريعة بين طلاب جامعة المجمعة. حيث كان أكثر من ربع المشاركين يستهلكون وبشكل يومي الوجبات السريعة وتعد هذه النسبة مرتفعة ومثيرة للقلق. لذلك ينبغي معالجة ذلك وزيادة الوعي لدى طلاب الجامعة لتقليل الآثار الصحية الضارة الناجمة عن استهلاك الوجبات السريعة.

**Introduction:** The term “fast food” was recognized in a dictionary by Merriam–Webster in 1951.<sup>1</sup> Fast food is a term given to food that can be prepared and served quickly. Usually it is a term given to junk food or food preheated or precooked. It is one of the largest influences on

the community as a whole especially on children, teens, and even adults.<sup>2</sup> It is believed that people who consume fast food on daily basis lacks good nutrition and physical condition. Fast food is definitely a problem that concerns us as a society.<sup>3</sup> The fast food market

in Saudi Arabia is expected to reach \$4.5 billion in gross sales by 2015, driven by growing demand from its population,<sup>4</sup> while obesity considered as a risk factor for many disease, consumption of fast food are reasons for increasing prevalence of obesity.<sup>5</sup> One study in United States reported that during 2007–2010, adults consumed, on average, 11.3% of their total daily calories from fast food; the highest percentage of calories from fast food was consumed among adults who were aged 20–39.<sup>6</sup> While in Saudi Arabia there are not clear statistics about consumption of fast food, but one news indicate that the rate of consumption of fast food among Saudi families reached 85% daily.<sup>7</sup> Available studies in Eastern Mediterranean countries indicate that fast food related disease like obesity has reached at an alarming level among both children and adults.<sup>8</sup> Another study conducted in Karachi, Pakistan, revealed that 58.3% students were of normal weight and 41.7% were overweight. In-parallel with the rapidly developing technology, eating habits also undergo changes. Although nutrition is important for all segments of the society, it is of a different importance for university students.<sup>9</sup> Individuals, who gain independence in this period, start to decide on their eating preferences, to eat out more frequently and to get influenced by their circle of friends more. Therefore, they tend to consume those foods that are deemed unhealthy such as fizzy drinks and fast-food more. Fast-food has become a significant symbol for the modern culture as it

satisfies people in a relatively short time.<sup>10</sup> College students are highly exposed to unhealthy eating habits leading to body weight gain.<sup>11</sup> According to WHO, obesity is generally more common among women than men.<sup>12</sup> Therefore, we planned this study to determine the prevalence of fast food consumption among Colleges' students; to associate gender with factors that leads to fast food consumption and to measure the level of awareness in association with gender about fast food consumption effects in students studying at Almajmaah University.

### **Methodology**

It was an observational cross-sectional study conducted in Majmaah University, the main campus is located in Majmaah City and has branches in other cities like, Alzulfi, Alghat, Hotatsudair, Ushirat Sudair, Rumah, and others. The study was conducted from December 2013 - May 2014. The target population was students of either gender studying in Majmaah University. A total of 505 students (325) males and (180) females aged between 18-26 years were randomly chosen from Colleges of Almajmaah University. The data was collected by Simple random sampling, using computer generated list Using direct investigation method. The questionnaire was adapted from study of "Consumption of Fast Food among the students of Faculty Technology at University Malaysia Pahang" and was validated by pilot study. Prior to filling out the questionnaire, the students were informed about the study and were given

instructions about how to fill out the questionnaire completely and truthfully. Ethical approval was also sought from Department of Essential and Health Sciences Research Center at Majmaah University. The data entered and analyzed by using SPSS 22.0. Mean $\pm$ S.D is reported for quantitative variables like age etc. Frequencies and percentages are reported for qualitative variables. Pearson Chi-square and Fisher Exact test were applied to observe associations between qualitative variables. A p-value of  $<0.05$  was considered as statistically significant.

## Results

The prevalence of fast food consumption among Majmmah University students was calculated on basis of two parameters; as a usual meal and weekly consumption. Two hundred ten participants (41.6%) choose fast food as their usual meal.

Fast food as a usual meal was consumed more by male students as compared to female student ( $p<0.001$ ). Majority of the students 430 (85.15%) were consuming fast food at least once a week. Two hundred and one (39.8%) of the participants (137 males, 64 females) were consuming fast food 2-3 times a week. Sixty (11.9%) of the participants (54 males, 6 females) were consuming fast food 4-5 times a week. While 65 (12.9%) of the participants (55 male, 10 female) were consuming fast food more than 5 times per week.

*Table1: Fast food consumption among affected population.*

Type of disease	Yes	No
Obesity	395 (78.2%)	192(38%)
Hypercholesterolemia	183 (36.2%)	322 (63.8%)
Heart Disease	137 (27.1%)	368 (72.9%)
Diabetes	131 (25.9%)	374 (74.1%)
Hypertension	88 (17.4%)	417 (82.6%)
Gallstones	35 (6.9%)	470 (93.1%)
Arthritis	30 (5.9%)	475 (94.1%)
Sleep Apnea	24 (4.8%)	481 (95.2%)
Others	19 (3.8%)	486 (96.2%)

Weekly consumption of fast food was more by male students as compared to female students ( $p<0.001$ ). About factors that leads to fast food consumption, we asked questions about: Distances of restaurants, taste, advertisements, lack of cooking skills, time limitation, prices, friends, and staying away of family. Significant association was observed between gender of participants and taste ( $p<0.001$ ), showing that female students were attracted by taste more than male students.

Female students were attracted to advertisements more than male students ( $p<0.001$ ). Consuming fast food and staying away from the family was significantly more in male students as compared to female students ( $p<0.001$ ). We also tried to measure the level of awareness among our participants regarding fast food consumption. About knowing of diseases

that fast food may cause 273 (54.1%) of participants knew about it.

*Table 2: Major factors affecting fast food consumption.*

Factor	Yes	No
Are Restaurants near?	313 (62%)	192(38%)
Restaurants Distance Effect	304 (60.2%)	201 (39.8%)
Taste	218 (43.2%)	287 (56.8%)
Friends	168 (33.3%)	337 (66.7%)
Time Limitations	151 (29.9%)	354 (70.1%)
Staying away of Family	119 (23.6%)	386 (76.4%)
Advertisements	86 (17%)	419 (83%)
Lacking of Cooking Skills	66 (13.1%)	439 (86.9%)
Prices	58 (11.5%)	447 (88.5%)

Majority of the students 395 (78.2%) responded that obesity is a major disease caused by consumption of fast food, followed by one quarter of the students who said hypercholesterolemia and about one fifth of the students said diabetes, hypertension and heart disease. Significant association was observed between gender of participants and knowing about diseases,  $p < 0.042$ , showing that female students knowledge about diseases that fast food may cause was more than male students.

No significant association was observed between Gender and knowledge about natural information and knowledge about ingredients on choosing fast food meals. About trying to lower or stop fast food consumption, 434 (85.9%) students answered “yes”, whereas, 68 (13.5%)

of the students said they will not stop or try to lower their fast food consumption.

## Discussion

Our study revealed that the prevalence of fast food consumption among Almajmaah university male and female students is high. In our study 85.15% of the students aged 18 to 26 years went to fast-food restaurants at least once per week. Our study is the first of its kind in the Kingdom which aims to determine prevalence of fast food consumption among important bridge of the community i.e. university students. A cross-sectional study conducted at Rass, Qassim University, Saudi Arabia<sup>14</sup> revealed that 21.8% of the students were overweight and 15.7% were obese with strong association to the dietary habits including fast food consumption, but it differs than our study as the prevalence in 85.15%. In another similar study<sup>15</sup> on frequent consumption of fast foods, low servings per day of fruits, vegetables, milk and dairy products, frequent consumption of sweets, candy and carbonated drinks was mild to moderate, but in our study prevalence is much higher. That may be because it was conducted on college students who are usually busy with educational activities with limited time, more responsibilities resulting in more exposure to fast food. Another study conducted in Dammam, Saudi Arabia<sup>16</sup> revealed that fast food rich in fat and calories from restaurants was popular among majority (98.2%) of the students. In our study, the prevalence is also high as it involves young

female students with almost same circumstances. Yet another study conducted on association between fast-food consumption and obesity in Michigan adults<sup>17</sup> showed that 80% of adults went to fast-food restaurants at least once per month which is as high prevalence as in our study because of similar age groups, educational activities and family issues. We compared our results related to reasons for eating fast food with a relatively similar study conducted in 2009;<sup>18</sup> the distance from the restaurant (60.2% Vs 71%), the taste (43.2% Vs 69%), sitting with friends (33.3% Vs 31%), limited time (29.9% Vs 46.8%) and being away from family (23.6% Vs 21%) advertisements (17% Vs 47%). In our study when gender differences were compared in relation to fast food consumption and nutrition self-assessments, female students' significantly strongly agreed with the supra stated statements than males, when this compared with a study<sup>18</sup> conducted in 2008 no significant association was observed. Regarding knowledge about the diseases that fast food disease may cause, females significantly knew about it more than the males ( $p>0.001$ ). When the classification of diseases were studied: fast food causes; obesity was reported by (78.2%) of the participants, hypercholesterolemia (36.2%), heart diseases (27.1%), diabetes (25.9%), hypertension (17.4%), gallstones (6.9%), arthritis (5.9%) and sleep apnea (4.8%). When these results were compared to a study conducted in Bangladesh<sup>18</sup> approximately 98% students were concerned about the negative

effects associated with fast food consumption, and 60% pointed out obesity and weight gain as the most prominent consequence of excessive fast food consumption. About trying to lower or stop fast food consumption a total of 85.9 % want to reduced, lower, or stop their fast food consumption. No similar study was found to compare our results with.

### **Recommendation**

We recommend to the fast food restaurants; to offer alternative meals with more healthy/natural components; awareness among the community should be developed related to diseases that fast food consumption causes, for this help from print or electronic media may be sought.

### **Conclusion**

The prevalence of fast food consumption among Almajmaah University students is very high. More than one quarter of the participants were consuming fast food daily which itself is alarming. Some intervention awareness programmed should be conducted to minimize the harmful effects of fast food consumption.

### **References**

1. Hopping BN, Erber E, Mead E, Sheehy T, Roache C, Sharma S. Socioeconomic indicators and frequency of traditional food, junk food, and fruit and vegetable consumption amongst Inuit adults in the Canadian Arctic. *J Hum Nutr Diet.* 2010; 23(Suppl 1):51–8.
2. Nicki R. Colledge, Brian R. Walker, Stuart H. Ralston, Davidson's Principles and Practice of Medicine. 21st ed. Elsevier: 2010; 5:117
3. Zeke Brynat. Who Consumes Fast-Food and Why? *SS-AAEA Journal of Agricultural Economics.* 2011. Vol 3, pp44-51.



4. Kumar P, Clark M, Fast food consumption in Middle East region. A step forward. Elsevier: 2009; 5,228.
5. Musaiger AO. Overweight and Obesity in Eastern Mediterranean Region: Prevalence and Possible Causes. Journal of Obesity. Volume 2011 (2011), Article ID 407237, 17 pages.
6. Shakila Ahmed, Imran Ali, Zabeen Jilani, Mrkran Usman. Fast-Food Consumption Habits of University Students: The Sample of Ankara Turkey. Pakistan Journal of Nutrition 2012; 11 (3): 265-269.
7. Huang TT, Harris KJ, Lee RE, Nazir N, Born W, Kaur H: Assessing Overweight, Obesity, Diet, and Physical Activity in College Students. World Health Organization: Obesity: preventing and managing the global epidemic. Report of a WHO consultation 2009.
8. Chang Pong, Jinla Buli, Mark Sam, Brandon macallum. Fast food for children and adolescents. Journal of Medical Sciences 2009;23(34), pp21 – 28.
9. Nighat Nisar, Majid Hafeez Qadri, Kiran Fatima. Dietary habits and life style among the students of a private Medical University Karachi, Pakistan, JPMA 2005;21(2):98-103.
10. Al-Rethaiaa AS, Fahmy AE, Al-Shwaiyat NM. Obesity and eating habits among college students in Saudi Arabia: a cross sectional study. Nutr J. 2010;9:39.
11. Amin TT, Al-Sultan AI, Ali A. Overweight and Obesity and their Association with Dietary Habits, and Sociodemographic Characteristics Among Male Primary School Children in Al-Hassa, Kingdom of Saudi Arabia. Indian J Community Med. 2008; 33(3):172–181.
12. Gunes FE, Bekiroglu N, Imeryuz N, Agirbasli M. Relation between eating habits and a high body mass index among freshman students: a cross-sectional study. J Am Coll Nutr. 2012;31(3):167-74.
13. Anderson B, Rafferty AP, Lyon-Callo S, Fussman C, Imes G. Fast-food consumption and obesity among Michigan adults. Prev Chronic Dis. 2011;8(4):A71. Epub 2011 Jun 15.
14. Rasheed P1, Al-Kunji AA, Al-Saffar BM, Al-Abdul Karim HM, Al-Thawadi MI. Are young arab women eating a healthy diet? A qualitative dietary study among college hostel students. J Family Community Med. 1999;6(2):17-22.
15. Akman M1, Akan H, Izbirak G, Tanriöver Ö, Tilev SM, Yildiz A, Tektaş S, Vitrinel A, Hayran O. Eating patterns of Turkish adolescents: a cross-sectional survey. Nutr J. 2010;19:9-67
16. Rydell SA1, Harnack LJ, Oakes JM, Story M, Jeffery RW, French SA. Why eat at fast-food restaurants: reported reasons among frequent consumers. J Am Diet Assoc. 2008;108(12):2066-70.
17. Morse KL1, Driskell JA. Observed sex differences in fast-food consumption and nutrition self-assessments and beliefs of college students. Nutr Res. 2009;29(3):173-9.
18. Munmun Shabnam Bipasha, Shatabdi Goon. Fast food preferences and food habits among students of private universities in Bangladesh. South East Asia Journal Of Public Health 2013;3(1):61-64.

**REVIEW ARTICLE****Athletic Pubalgia – A Review of Literature.****Qassim I. Muaidi\*<sup>1</sup>**\* For correspondence [qmuaidi@uod.edu.sa](mailto:qmuaidi@uod.edu.sa)

\*Vice Dean for Academic Affairs &amp; Chairman of Physical Therapy Department, College of Applied Medical Sciences, Dammam University

Submitted on: 15<sup>th</sup> September 2015, Accepted on: 11<sup>th</sup> October 2015**Abstract**

Athletic pubalgia is commonly seen in sports that require athletes to perform repetitive kicking, cutting, pivoting and sharp turns such as soccer, cricket, rugby, ice hockey etc. It is caused by immoderate and disproportionate application of shear forces through the common attachments (pelvic) of the abdominal muscles and hip adductors. At present, there is no unanimity of opinion about the constituents of this diagnosis. Since it is not easy to define a clear diagnosis that is based on physical examination, methods such as ultrasonography and MRI are commonly used to rule out the existence of other conditions. Though surgeries are better than traditional treatment, laparoscopic surgery has a shorter recovery time as compared to open surgeries. This article summarises the present data regarding anatomy, biomechanics, clinical presentation, differential diagnosis, conservative treatment, preoperative and post-operative rehabilitation of athletic pubalgia. The information obtained from such a research work will help in understanding pathogenesis in a better way, build-up evidence-oriented screening and check for better surgical methods and effectively manage both post-surgical and conservative rehabilitation.

**Key Words:** Abdominal Muscles, athletes, laparoscopy, physical examination, running.

**المخلص**

الفتق عند الرياضيين عادة ما يحصل في الألعاب الرياضية التي تتطلب أداء متكرر من الحركات مثل الركل وتغيير الاتجاهات والدوران بشكل حاد وسريع مثل لعبة كرة القدم، والركبي، والكريكت، ولعبة الهوكي على الجليد. وتكون ناتجة تعرض عضلات الحوض والفخذ الداخلية والبطن لقوة مفرطة وغير متناسبة مع قوة هذه العضلات. في الوقت الحاضر، لا يوجد إجماع في الرأي حول مكونات هذا التشخيص. ونظرا لأنه ليس من السهل تحديد تشخيص دقيق مبني فقط على الفحص السريري، فإن عدة طرق أخرى مثل الموجات فوق الصوتية والتصوير بالرنين المغناطيسي تستخدم لاستبعاد وجود إصابات أو امراض أخرى. على الرغم من ان العمليات الجراحية هي أفضل من العلاج التقليدي، الا ان وقت الشفاء لجراحة المناظير أقصر من العمليات الجراحية المفتوحة. هذا المقال يلخص البيانات الحالية المتعلقة بالتشريح، الميكانيكا الحيوية، الطرق السريرية، التشخيص التفريقي، العلاج الغير جراحي والجراحي، والتأهيل قبل وبعد الجراحة للفتق عند الرياضيين. ان هذه المعلومات التي تم الحصول عليها من الأبحاث العلمية ستساعد في فهم مصدر وتطور هذا المرض او الإصابة بشكل أفضل، وبناء ادلة مبنية على التشخيص والفحص الدقيق والتحقق من الأساليب الجراحية الأفضل، وفضل طرق التأهيل الغير جراحية والتأهيل بعد العمليات الجراحية.

**Introduction**

Athletic pubalgia, also known as Sports hernia or Gilmore's groin is commonly seen among the athletes who perform repetitive high speed kicking, turning or pivoting commonly seen in cricket, tennis, basket ball and most frequently soccer and hockey.<sup>1-5</sup> Athletic pubalgia refers to the chronic pain in the inguinal or pubic-area

which is present with exertion and not explainable preoperatively by a palpable hernia or other medical diagnosis.<sup>6</sup> It may be unilateral or bilateral depends up on the location of pain and there may be associated medial thigh symptoms related to the adductors of the hip.<sup>3,7</sup> It has been reported among 5% to 28% of athletes<sup>9</sup> and most commonly diagnosed among

male athletes.<sup>8</sup> Around 2% of female athletes have been diagnosed with athletic pubalgia.<sup>8</sup> This variation in the incidence of pubalgia may be because of the anatomical differences between male and female rather than participation.<sup>10</sup> Females have a wider pelvis which makes more effective force distribution to the lower body<sup>11</sup> and the strength variation also makes contribution to high incidence of pubalgia among males. Literature is filled with complicated and by and large conflicting information regarding the cause, medical presentation, diagnosis, and management of sports hernia in athletes.<sup>1,4</sup> It is important that the clinicians should have a thorough knowledge about the condition as there is a high chance of misdiagnosis.

**Applied Anatomy and Biomechanics:** Rectus abdominus makes a conjoined tendon with transversus abdominus muscle and insert at or near the pubis. Across from these two muscles, and directly opposing their forces, is that the adductor muscle longus. Rectus abdominis muscle originates from the upper part of the symphysis pubis, with distinction often able to be made between a lateral and a medial head on both sides. Inferiorly, the medial head blends with its contralateral fellow; however, superiorly the medial heads diverge and are separated by the linea alba. The rectus abdominis has a sheet like configuration,<sup>12</sup> with a minimum of three intramuscular tendinous intersections, one at the level of the umbilicus,

one at the xiphoid tip, and one between these two intersections. Sometimes, a fourth intersection may be present inferior to the umbilicus. The intersections extend through only to the anterior surface of the muscle, where they blend with the anterior rectus sheath. This latter structure covers the complete anterior aspect of each rectus abdominis and attaches onto the periosteum of the pubic bone anterior and adjacent to the rectus abdominis origin. The adductor muscle longus and adductor muscle brevis possess an extensive insertion on the femur. Combined with the insertion of the gracilis onto the tibia, these three tendons converge superiorly and obtain an origin close to each other on a narrow portion of the pubic body just lateral to the symphysis.

The adductor muscle tendon has its origin almost directly in line with the origin of the more superiorly placed tendon of the rectus abdominis, with the superficial fibers of these two tendons in direct continuity, coursing over the pubic crest. Farther posteriorly and slightly laterally lies the origin of the adductor muscle brevis. Some of the medial fibers of the adductor muscle longus and adductor muscle brevis tendons attach directly onto the symphyseal capsular tissues and intra articular disk.<sup>13</sup> When an athlete suffers a sports hernia, these opposing forces (upward pull from conjoined tendon and downward pull from adductor longus) cause disruption of the muscles at their insertion site (pubis), leading to groin pain. What happens is

that in repetitive kicking, pivoting or turning at high speeds the conjoined tendon exerts upward force and rotates the trunk, and the adductor muscle longus exert downward force and rotate the upper leg. The application of these excessive amounts of force in an imbalanced manner causes Athletic pubalgia. Often, the core muscles are less developed than the lower extremity muscles and disruption disproportionately involves the conjoined tendon leading to athletic pubalgia.

*Signs and Symptoms:* Athletic pubalgia is commonly seen among males than females. Most of the time the onset is insidious and in some cases an initial sudden tearing sensation may be felt. Pain is usually exercise related and felt on the lower side of the rectus abdominis muscle and it may radiate toward the testicular area, supra pubic region or adductor region. The pain may get aggravated by a sudden acceleration, cutting or even cough or sneezing. There may associate stiffness along with pain and it may persist for more than two to four days after competition. Localized tenderness at or above pubic crest of the effected side and pain will be present during resisted sit-ups. Resisted hip abduction will be painful and a painful squeeze test in supine position or 90° hip flexion position.

*Diagnosis:* The following criteria required for a diagnosis of athletic pubalgia includes: a clinical setting of chronic groin pain which develops during exercise, is felt over the lower

lateral side of the rectus abdominis muscle with or without radiating to the testis or medial thigh muscles, and is aggravated by sudden acceleration, cutting and kicking, pivoting, sit-ups and coughing or sneezing; Subtle but consistent physical examination findings (Physical examination will not find an actual hernia, even though most of these signs and symptoms lead health care professionals to believe that one is present).<sup>10,17-21</sup> The Valsalva maneuver, along with muscle testing of the hip adductors, rectus abdominis, rectus femoris, and iliopsoas, will produce some discomfort.<sup>20</sup> Differential diagnosis has to be done to rule other causes of groin pain and appropriate imaging features. All of these criteria must be simultaneously present, because: (1) there are numerous other potential causes for groin pain in athletes; and (2) asymptomatic direct inguinal hernias are common in the general population.<sup>22</sup>

*Differential diagnosis:* Adductor Longus Tendinosis and Tenoperiostitis: It usually occurs with persistent overuse of the adductor longus muscle.<sup>23</sup> Adductor tendinosis is generally related to a rectus abdominis–adductor muscle longus aponeurotic injury. However, these type of injuries can be treated conservative management. The most common area of injury is proximal myotendinous junction. Pain and discomfort is in the upper medial thigh may get worsened by a resisted adduction of the hip and a tenderness can be felt on the proximal part of the tendon. The MRI shows thickening of the

adductor muscle longus tendon on the affected area, intra-tendinous tear, calcification, tendon avulsion, or a higher-grade and more distal adductor strain.

*Osteitis Pubis:* It is believed to result from instability of the pubic symphysis because of chronic repetitive shear and distraction injuries and unbalanced tensile stress from the muscle attachments of the pubic symphysis. The biomechanical changes leads to an inflammatory response, with osteitis and periostitis. Onset of pain over the pubis that may cause referred pain in the suprapubic region, adductor origin, and groin. Radiographs of the pubic symphysis may show subchondral sclerosis, symphyseal irregularity, and bone resorption.<sup>24</sup> MR images show diffuse marrow edema extending from the subchondral plate and often involving both pubic bodies.<sup>25,26</sup> In addition, periostitis, erosions, irregularity in articular surface, anterior and posterior osteophytes, and subchondral cysts may also be seen on MRI.<sup>27</sup> The myotendinous attachments are preserved unless there is coexistent tendinopathy.<sup>28</sup> The marrow edema in an osteitis pubis extends across the entire anteroposterior dimension of the pubic body.

*Hockey Goalie–Baseball Pitcher Syndrome:* Groin pain in this unusual syndrome result from an epimysial or myofascial herniation of the adductor longus muscle belly several centimetres away from the site of its pubic attachment.<sup>29</sup> A number of authors have

suggested a link with chronic repetitive stress at sites of neurovascular penetration.<sup>30</sup> Acute onset of pain is usually experienced by patients in this condition, which may be persistent or may intermittently increase after stretching. Physical examination often leads to the identification of a site of pain distal to the symphysis pubis, over the herniated muscle. Imaging findings shows edema of the adductor longus muscle belly, which can possibly occupy a central position in the herniated area, and a focal muscle bulge indicative of herniation.<sup>31</sup>

*Pubic Stress Fracture:* Fatigue fractures are usually seen in athletes, especially in long-distance runners who are supposed to reflect abnormal chronic repetitive stress that surpass the bone's capability to remodel itself. Inferior pubic ramus is the most common site of pelvic stress fracture; however, stress fractures of the femoral neck are more prevalent and may cause groin pain.<sup>32</sup> Risk factors for stress fracture include female sex, malnutrition, and changes in the training program that place a greater amount of stress on the bone. Patients usually present with a gradual onset of pain, which, at an initial phase of injury, is aggravated by exercise and relieved by rest. If pain persists even during rest, it indicates a more advanced phase of injury. On palpation, tenderness is usually present over pubic ramus.<sup>33</sup> MRI finding shows areas of marrow edema obtained with fluid-sensitive sequences. There is also a possibility that periosteal edema or calluses and edema in the



adjacent muscles and soft tissues are present in some cases.

*Septic Arthritis–Osteomyelitis:* Infection of the pubic symphysis is commonly seen in postpartum female patients, patients with pelvic malignancies and patients subjected to direct instrumentation of the pubic symphysis, but it is also seen in athletes without such risk factors.<sup>34</sup> The infection is most often caused by *Staphylococcus aureus* and may result from a hematogenous spread rather than direct inoculation. Patients usually present with fever and an acute onset of pubic pain that can radiate up to the groin. The radiographic and MRI features of pubic septic arthritis–osteomyelitis and osteitis pubis are identical, and clinicians may depend on the duration of symptoms and the presence of risk factors to help them distinguish between these possibilities. If the diagnosis is not definite, a biopsy test is desired.

*Acetabular Labral Tear:* There is complete overlap between the clinical manifestations of athletic pubalgia and the pathologic conditions associated with the hip. Some of the hip disorders which may result in groin pain are osteoarthritis, synovitis, intra-articular bodies, and acetabular labral tears. The ant-superior part of the labrum is quite vulnerable to injuries, particularly during external rotation and hyperextension.<sup>35</sup> The labrum is comparatively poorly vascularized and its anterior-superior aspect is actually considered to be particularly weak. Labral injuries predispose the hip to paced

articular cartilage loss, especially near to the tear, a condition that is considered to be the cause of alterations and instability in contact forces of the articular cartilages.<sup>34</sup> Abnormal morphology and signal intensity of the labrum at routine MR imaging of the hip are indicative of the diagnosis. Paralabral cysts linked with acetabular labral tears can be clearly seen on the images obtained with fluid-sensitive sequences. Additionally, MR imaging plays its role in determining whether the morphologic features of the femoral head and acetabulum predispose a patient to labral tears and femoroacetabular impingement.

*Internal Snapping Hip (Coxa Saltans):* Internal snapping hip is a rare cause of pain in the inguinal region and anterior part of the hip. The musculo tendentious part of the iliopsoas muscle snaps with head and anterior capsule of the femur causing an obstruction to the movement. This chronic repetitive motion may leads to tendinosis and iliopsoas bursitis. Magnetic Resonance Imaging of the hip in a patient with internal snapping may seem to be normal, but iliopsoas bursitis is frequently present.<sup>36,37</sup> Intermediate intra substance signal intensity and Iliopsoas tendon thickening are rare findings, but when they are observed over the spinal process or iliopectineal eminence, they are indicative of the diagnosis. Lastly, an osseous spur budding from the iliopectineal eminence may predispose the patient to growing internal snapping. Ultrasonography is often helpful for

visualizing the tendon as it passes over the iliopectineal eminence during dynamic maneuvers in addition for detecting bursitis.

*Osteoid Osteoma:* Osteoid osteoma is a benign bone-forming tumor generally observed in the patients between the ages of 5-30 years. It is generally occurring in the long bones, specifically the tibia and femur and in the phalanges of the toes and fingers. Lesions that include the pubic bones may repeat the symptoms of athletic pubalgia. Osteoid osteomas may appear in medullary, cortical, subcortical or periosteal sites. They typically occur as a central radiolucent nidus with a varying degree of calcification, surrounded by different zones of cortical thickening in addition to non aggressive periosteal reaction. Intramedullary lesions as well as lesions within a joint capsule are often accompanied by a small amount of or no surrounding sclerosis due to the lack of contiguity with the nearest periosteum. CT and Radiography are generally diagnostic for osteoid osteoma and may be used to give direction to biopsy and treatments like percutaneous ablation.<sup>38</sup>

*Nerve Entrapment Syndromes:* Several nerves, including the femoral, obturator, iliohypogastric, genitofemoral, lateral femoral cutaneous nerves and ilioinguinal provide motor and sensory innervation to the upper thighs and groin. Entrapment of any of these structures may result in groin pain mimicking athletic pubalgia. Patients with obturator nerve entrapment,

mostly present along with aching pain near the adductor origin which may radiate to the knee and also worsen during the exercise. The nerve entrapment has been attributed to fascial thickening of the adductor compartment, specifically along the anterior aspect of the adductor brevis.<sup>39,40</sup> Other possible causes of obturator nerve entrapment involve a mass effect from an obturator hernia a pelvic fracture, an obturator hernia and an acetabular paralabral cyst.<sup>41,42</sup> Femoral nerve entrapment has been diagnosed in patients after surgical procedures such as herniorrhaphy, hip arthroplasty, abdominal hysterectomy, entrapment of the genitofemoral and ilioinguinal nerves may be seen after abdominal surgery in muscle hypertrophy or in blunt trauma.<sup>43</sup> Although the nerves in this area are difficult to detect at regular MRI, it may be helpful for diagnosing entrapment. MRI findings of muscle denervation-related edema or atrophy with a characteristic pattern may be suggestive of entrapment of a specific nerve.

*Apophysitis:* There are several pelvic apophyses which may be injured by forceful or repetitive muscle contractions during athletic activities before complete skeletal maturity is achieved i.e. between the ages of 12 and 22 years. Groin pain mimicking athletic pubalgia may be present in inflammation of the apophyses of the anterior-inferior iliac spine, where actually the rectus femoris originates and the anterior-superior iliac spine at the originating point of the

Sartorius and the tensor fascia lata. Athletes who participate in sports involving frequent kicking have a tendency to develop painful apophysitis in such locations.

*Urological disease:* Other commonly seen causes of obscure groin pain include prostatitis,<sup>44</sup> urethritis, epididymitis and hydrocele.<sup>45</sup>

*Connective tissue disease:* Various connective tissue disease, such as rheumatoid arthritis should be ruled out. Diseases such as gout, ankylosing spondylitis, Reiter's syndrome and other spondylarthritides may also need to be investigated.<sup>46</sup>

*Spinal and hip abnormalities:* Early assessment of the spine and hip has to be done to identify sacrolumbar abnormalities and sinister lesions such as bone tumours. Old osteochondritis of the vertebral bodies, disc lesions at L1 or L2, and crush fractures can cause radicular pain and imitate athletic pubalgia.

*Imaging:* Many authors suggest that Magnetic Resonance Imaging (MRI) is having a limited ability to identify an athletic pubalgia but it can be used to rule out any of the alternative diagnoses; nevertheless others have prompted that Athletic pubalgia almost always show abnormalities in MRI.<sup>47-51</sup> The two clearly seen patterns of injury involve lateral rectus abdominis/adductor aponeurotic injury just near to the midline rectus abdominis/adductor aponeurotic plate injury<sup>7</sup> and external inguinal ring. Although some

degree of pubic marrow enhancement is generally reported, these findings do not foretell which patients will gain from the surgery.<sup>7</sup>

*Management:* The initial stage of treatment for any kind of muscle injury generally includes rest, ice and proper compression. Resting is obviously needed to avoid any further deterioration or aggravation of the condition. Resting period can be anywhere from few weeks to months depending upon the condition's severity and application of ice to the injured area for about 10-15 minutes every four hours all over the day help in giving some relief in the swelling and pain caused by injury in Athletic pubalgia. Compression shorts and warm pants also play their role in controlling the symptoms of athletes who still have some time for surgical intervention. However, these may not decrease the symptoms of Athletic pubalgia, but certainly help in curbing worsening of tear until the time of surgical intervention. Anti-inflammatory medications like ibuprofen may also be consumed to reduce the swelling and pain.

*Physiotherapy Management:* The most important aim of physical therapy management is to reduce the time lost to the player due to injury and enhance the availability of the player to the team as soon as possible. Pre-season screening of the athlete especially flexibility, posture and gait is having a major role in prevention of these type of injuries. Once the assessment is done athlete should involve a rigorous training which is specific to each

individual. Early part of functional rehabilitation should be taught to the athlete in advance to the surgery so that the athlete should have an early understanding regarding the exercises following surgery.

More emphasis should be given on the stabilising role of the transverses abdominis, which is always neglected, internal oblique, gluteal muscles and adductors of the hip. In the late preoperative/early postoperative phase, more stress should be given in maintaining the neutral position of the spine. This requires time and a detailed explanation of what is needed during what appear to be quite simple exercises for the 'ill-informed' patient. It is important that the rehabilitation program should be sports specific or occupation specific. The medical team should have an excellent understanding regarding the demands of the sport/occupation.<sup>52</sup> Appropriate preoperative and postoperative care must allow a full return to activity within just four to eight weeks from the date of surgery. A good coordination between the treating clinicians and the physiotherapy team for the preoperative and postoperative rehabilitation is recommended.

*Preoperative rehabilitation:* Preoperative rehabilitation involves; postural, range of motion and strength assessment, strengthening of gluteal muscles, transvers abdominis, erector spinae and abdominal muscles, educating the patient regarding post-operative rehabilitation,

core muscle strengthening by the use Swiss ball, medicine ball, pulleys etc.

*Surgery:* Once the non surgical management is found ineffective, surgery is a reasonable option. Athletic pubalgia surgery can be done with all three types (local anaesthesia, general or regional anaesthesia). Athletic pubalgia surgery centres on stabilizing the insertion of the conjoined tendon at the pubic tubercle. Additionally, superficial nerve is taken away because this nerve can get caught up with post-surgical scar and mimics the symptoms of Athletic pubalgia. In order to strengthen the inguinal floor a supportive mesh will be placed over the repair and extend through the inguinal canal. The incision will be around four to five centimetre length. In order to decrease post-surgical discomfort local anaesthetics will be given before closing of the incision. The initial Athletic pubalgia surgery does not take care of the discomfort at the adductor insertion site. If the conjoined tendon is sufficiently supported by the Athletic pubalgia surgery, adductor discomfort is can be treated in postoperative rehabilitation. Rarely does the adductor needs operative release. Other surgical techniques involves an altered Bassini repair, in this technique the transversal fascia from the deep to superficial ring was released. Intra operatively, all athletes were diagnosed for thinning and a variable degree of bulging of the posterior inguinal wall. The conjoined tendon used to be attached to the inguinal ligament in a double-

layer continuous suture, as against the interrupted original Bassini repair; with a running 2/0 prolene suture.<sup>53</sup> A concurrent percutaneous adductor release was done by an orthopaedic surgeon in all the cases. A bilateral tenotomy was done with complete transaction of the epimysial fibres of the adductor longus tendon which is two centimetre from the pubic insertion. For those who are having persistent adductor symptoms even after post surgical rehabilitation or for those whose major complain was related to adductor muscle, non-operative options do exist. Adductor problems, that are not responsive to physical therapy can be treated with Platelet-Rich Plasma injection, which is an office-based procedure. Post-procedure rehabilitation will be continued, and it is rare that a second injection is needed. This technique has replaced operative release of the adductor insertion.

*Post Surgical Rehabilitation:* Postoperative rehabilitation can be started with isometric exercises of abdominal muscles and muscles around the hip joint. Early spinal mobilization program can also be initiated. Increase walking using time as limiting factor, increasing by five minutes per day can be initiated from 2<sup>nd</sup> week onwards and active assisted exercises and isokinetic program in functional standing position can be incorporated at the end of week number.<sup>2</sup> Functional rehabilitation program by using gymnastic ball, mobility and stability exercises and the exercise to improve

cardiovascular system can be started with from 3<sup>rd</sup> week onwards. Early sports/occupational specific rehabilitation program can be incorporated with the existing rehabilitation program in the 4<sup>th</sup> and 5<sup>th</sup> weeks of surgery. The person can return to normal activities/sports by 5<sup>th</sup> or 6<sup>th</sup> week according to the functional reassessment.

### **Conclusion**

Timely recognition is crucial in case of injury diagnosis, which can be done through thorough examination of the injury. This may include palpation of the areas of concern and manual muscle testing to exclude the other possible conditions. Mostly, the patient will complain about chronic pain in the regions such as lower abdomen, pelvic region and thighs, which all radiates to the groin region. If a clinician encounters such complaints/symptoms, then, he/she should examine the patient for Athletic pubalgia. However, one needs to be aware about the fact that these findings will not confirm the diagnosis of Athletic pubalgia, so clinicians should have a thorough knowledge of the other conditions which may mimic the symptoms. Most imaging studies are only useful to exclude other diagnoses. Patient will typically not respond to conservative/non operative treatment, so we can switch over to surgical treatment if the patient does not respond to non operative treatment even after six weeks. Usually patient returns to his pre injury level



within 6 weeks after laparoscopic repair and within 1-6 months in case of an open repair.

### Acknowledgment

The author acknowledge the immense help received from the faculty members of Department of Physical therapy, University of Dammam for the completion of the review.

### References

1. Kachingwe AF, Grech S. Proposed algorithm for the management of athletes with athletic pubalgia (sports hernia): a case series. *The Journal of orthopaedic and sports physical therapy* 2008; 38: 768-81.
2. Holmich P. Long-standing groin pain in sportspeople falls into three primary patterns, a "clinical entity" approach: a prospective study of 207 patients. *British journal of sports medicine* 2007; 41: 247-52; discussion 52.
3. Meyers WC, Foley DP, Garrett WE, Lohnes JH, Mandlebaum BR. Management of severe lower abdominal or inguinal pain in high-performance athletes. PAIN (Performing Athletes with Abdominal or Inguinal Neuromuscular Pain Study Group). *The American journal of sports medicine* 2000; 28: 2-8.
4. Swan KG, Jr., Wolcott M. The athletic hernia: a systematic review. *Clinical orthopaedics and related research* 2007; 455:78-87.
5. Gilmore J. Groin pain in the soccer athlete: fact, fiction, and treatment. *Clinics in sports medicine* 1998; 17: 787-93
6. Hackney RG. The sports hernia: a cause of chronic groin pain. *British journal of sports medicine* 1993; 27: 58-62.
7. Preskitt JT. Sports hernia: the experience of Baylor University Medical Center at Dallas. *Proceedings (Baylor University Medical Center)* 2011; 24: 89.
8. Nam A, Brody F. Management and therapy for sports hernia. *Journal of the American College of surgeons* 2008; 206: 154-64.
9. Larson CM, Lohnes JH. Surgical management of athleticpubalgia. *Operative Techniques in Sports Medicine* 2002; 10: 228-32.
10. Diesen DL, Pappas TN. Sports hernias. *Advances in surgery* 2007; 41: 177-87.
11. Meyers WC, Yoo E, Devon ON, Jain N, Horner M, Lauencin C, et al. Understanding "sports hernia"(athletic pubalgia): the anatomic and pathophysiologic basis for abdominal and groin pain in athletes. *Operative techniques in sports medicine* 2012; 20: 33-45.
12. Rizk NN. A new description of the anterior abdominal wall in man and mammals. *Journal of anatomy* 1980; 131: 373.
13. Robinson P, Salehi F, Grainger A, Clemence M, Schilders E, O'Connor P, et al. Cadaveric and MRI study of the musculotendinous contributions to the capsule of the symphysis pubis. *American Journal of Roentgenology* 2007; 188: W440-W5.
14. Anderson K, Strickland SM, Warren R. Hip and groin injuries in athletes. *The American journal of sports medicine* 2001; 29: 521-33.
15. Brukner P, Khan K. *Clinical sports medicine*. McGraw Hill; 2006.
16. Joesting DR. Diagnosis and treatment of sportsman's hernia. *Current sports medicine reports* 2002; 1: 121-4.
17. Geisler P, Kelly E. Groin pains. *Training and Conditioning* 2008; 18: 31-5.
18. Schuricht A, Haut E, Wetzler M. Surgical options in the treatmentof sports hernia. *Operative Techniques in Sports Medicine* 2002; 10: 224-7.
19. Ahumada LA, Ashruf S, Espinosa-de-los-Monteros A, Long JN, Jorge I, Garth WP, et al. Athletic pubalgia: definition and surgical treatment. *Annals of plastic surgery* 2005; 55: 393-6.
20. Farber AJ, Wilckens JH. Sports hernia: diagnosis and therapeutic approach. *Journal of the American Academy of Orthopaedic Surgeons* 2007; 15: 507-14.
21. Paluska SA. An overview of hip injuries in running. *Sports medicine* 2005; 35: 991-1014.
22. Skandalakis JE, Gray SW, Skandalakis LJ, Colborn GL, Pemberton LB. Surgical anatomy of the inguinal area. *World journal of surgery* 1989; 13: 490-8.
23. Akermark C, Johansson C. Tenotomy of the adductor longus tendon in the treatment of chronic groin pain in athletes. *The American journal of sports medicine* 1992; 20: 640-3.
24. Vix Va, Ryu Cy. The adult symphysis pubis: normal and abnormal. *American Journal of Roentgenology* 1971; 112: 517-25.
25. Gibbon W, Hession P. Diseases of the pubis and pubic symphysis: MR imaging appearances. *AJR American journal of roentgenology* 1997; 169: 849-53.
26. Verrall G, Slavotinek J, Fon G. Incidence of pubic bone marrow oedema in Australian rules football players: relation to groin pain. *British journal of sports medicine* 2001; 35: 28-33.

27. Kunduracioglu B, Yilmaz C, Yorubulut M, Kudas S. Magnetic resonance findings of osteitis pubis. *Journal of Magnetic Resonance Imaging* 2007; 25: 535-9.
28. Brennan D, O'Connell MJ, Ryan M, Cunningham P, Taylor D, Cronin C, et al. Secondary Cleft Sign as a Marker of Injury in Athletes with Groin Pain: MR Image Appearance and Interpretation 1. *Radiology* 2005; 235: 162-7.
29. Meyers WC, Lanfranco A, Castellanos A. Surgical management of chronic lower abdominal and groin pain in high-performance athletes. *Current sports medicine reports* 2002; 1: 301-5.
30. Gokhale S. Three-dimensional sonography of muscle hernias. *Journal of ultrasound in medicine* 2007; 26: 239-42.
31. Mellado J, del Palomar LP. Muscle hernias of the lower leg: MRI findings. *Skeletal radiology* 1999; 28: 465-9.
32. LeBlanc KE, LeBlanc KA. Groin pain in athletes. *Hernia : the journal of hernias and abdominal wall surgery* 2003; 7: 68-71.
33. Lynch SA, Renström PA. Groin injuries in sport. *Sports medicine* 1999; 28: 137-44.
34. Morelli V, Espinoza L. Groin injuries and groin pain in athletes: part 2. Primary Care: Clinics in Office Practice 2005; 32: 185-200.
35. Overdeck KH, Palmer WE. Imaging of hip and groin injuries in athletes. In: *Seminars in musculoskeletal radiology*. 2004; Copyright© 2004 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.; p.41-55.
36. Brittenden J, Robinson P. Imaging of pelvic injuries in athletes. *Imaging* 2005; 78.
37. Blankenbaker DG, Tuite MJ. The painful hip: new concepts. *Skeletal radiology* 2006; 35: 352-70.
38. Cantwell CP, Obyrne J, Eustace S. Current trends in treatment of osteoid osteoma with an emphasis on radiofrequency ablation. *European radiology* 2004; 14: 607-17.
39. Harvey G, Bell S. Obturator neuropathy: an anatomic perspective. *Clinical orthopaedics and related research* 1999; 363: 203-11.
40. Bradshaw C, McCrory P, Bell S, Brukner P. Obturator Nerve Entrapment A Cause of Groin Pain in Athletes. *The American journal of sports medicine* 1997; 25: 402-8.
41. Modelli M, Giannini F, Guazzi G, Corbelli P. Obturator neuropathy due to obturator hernia. *Muscle & nerve* 2002; 26: 291-2.
42. Yukata K, Arai K, Yoshizumi Y, Tamano K, Imada K, Nakaima N. Obturator neuropathy caused by an acetabular labral cyst: MRI findings. *American Journal of Roentgenology* 2005; 184: S112-S4.
43. Morelli V, Weaver V. Groin injuries and groin pain in athletes: part 1. Primary Care: Clinics in Office Practice 2005; 32: 163-83.
44. Smedberg S, Broome A, Elmer O, Gullmo A. Herniography in the diagnosis of obscure groin pain. *Acta chirurgica scandinavica* 1984; 151: 663-7.
45. Taylor DC, Meyers WC, Moylan JA, Lohnes J, Bassett FH, Garrett WE. Abdominal musculature abnormalities as a cause of groin pain in athletes Inguinal hernias and pubalgia. *The American journal of sports medicine* 1991; 19: 239-42.
46. NIEPEL GA, Sitaj S. Enthesopathy. *Clinics in Rheumatic Diseases* 1979; 5: 857-72.
47. Lacroix VJ, Kinnear DG, Mulder DS, Brown RA. Lower abdominal pain syndrome in national hockey league players: a report of 11 cases. *Clinical Journal of Sport Medicine* 1998; 8: 5-9.
48. Nelson EN, Kassarian A, Palmer WE. MR imaging of sports-related groin pain. *Magnetic resonance imaging clinics of North America* 2005; 13: 727-42.
49. Van Veen R, de Baat P, Heijboer M, Kazemier G, Punt B, Dwarkasing R, et al. Successful endoscopic treatment of chronic groin pain in athletes. *Surgical endoscopy* 2007; 21: 189-93.
50. Kluin J, den Hoed PT, van Linschoten R, IJzerman JC, van Steensel CJ. Endoscopic evaluation and treatment of groin pain in the athlete. *The American journal of sports medicine* 2004; 32: 944-9.
51. Shortt CP, Zoga AC, Kavanagh EC, Meyers WC. Anatomy, pathology, and MRI findings in the sports hernia. In: *Seminars in musculoskeletal radiology*. 2008; © Thieme Medical Publishers; p.054-61.
52. Macintyre J, Johnson C, Schroeder EL. Groin pain in athletes. *Current sports medicine reports* 2006;5: 293-9.
53. Van Der Donckt K, Steenbrugge F, Van Den Abbeele K, Verdonk R, Verhelst M. Bassini's hernial repair and adductor longus tenotomy in the treatment of chronic groin pain in athletes. *Pain* 2003; 36: 38.

**VIEWPOINT**

## Teaching Biostatistics to Medical Undergraduates in Integrated Curriculum

Waqas Sami<sup>\*1</sup>, Mohammed AlMansour<sup>2</sup>, Tayyaba Waqas<sup>3</sup>

\* For correspondence [w.mahmood@mu.edu.sa](mailto:w.mahmood@mu.edu.sa)

<sup>1</sup>Waqas Sami, Lecturer of Biostatistics, Department of Public Health & Community Medicine, College of Medicine, Majmaah University, Almajmaah, Saudi Arabia.

<sup>2</sup>Assistant Professor, Family Medicine, Department of Public Health & Community Medicine, College of Medicine, Majmaah University, Almajmaah, Saudi Arabia.

<sup>3</sup>Biostatistician, Department of Continuous Medical Education, Punjab Institute of Cardiology, Jail Road, Lahore, Pakistan.

Submitted on: 4<sup>th</sup> January 2015, Accepted on: 6<sup>th</sup> September 2015

### Abstract

As medicine has moved to become more evidence based, biostatistics has become ever more important in practicing medicine and for the education of tomorrow's doctors. The role of biostatistics in medical education is now well recognized, and the curricula in almost all the medical colleges and postgraduate institutions across globe has provision of teaching biostatistics. In an integrated curriculum biostatistics and research methods, course should be inducted in-combination from 1st year of study and its Curricula should be developed by content experts with taking care of the spiral approach. Integration is now considered the best way to teach so the tutors should get maximum benefit of teaching biostatistics under this system. The curriculum contents should be updated at the end of course every time based on feedbacks and recent advancements in the field. Computer based approach should be used to make the learning more interactive. Assessment must be aligned with teaching methods, curriculum outcomes and course objectives.

**Key Words:** Biostatistics, Research Methods, Integrated Curriculum, Computer Based Approach

### الملخص

بما أنه كما هو معلوم بأن الطب أصبح يعتمد على البراهين، فقد أصبح الإحصاء الحيوي أهم من أي وقت مضى في ممارسة الطب وتعليم أطباء الغد. ومن المسلم به الآن دور الإحصاء الحيوي في التعليم الطبي والمناهج الدراسية تقريبا في جميع الكليات الطبية والمؤسسات التي تمنح دراسات عليا في الطب في جميع أنحاء العالم. في المناهج المتكاملة، مقرر الإحصاء الحيوي ومناهج البحث، ينبغي أن يدخل في مناهج السنة الأولى للدراسة والمناهج ينبغي أن توضع من قبل خبراء التخصص مع الأخذ بالاعتبار المقاربة الدائمة للمنهج الكلي للكلية. يعتبر التكامل الآن أفضل وسيلة للتعليم وبالتالي فإن المعلمين يجب ان يحصلوا على الفائدة القصوى من تدريس الإحصاء الحيوي في ظل هذا النظام. يجب أن يتم تحديث محتويات المنهج الدراسي في نهاية كل مقرر على أساس التغذية الراجعة والتطورات الأخيرة في هذا المجال. وينبغي استخدام الكمبيوتر لجعل التعلم أكثر تفاعلية. التقييم يجب أن يكون متوافقا مع طرق التدريس ومخرجات المنهج وأهداف المقرر.

### Introduction

Prior to the twentieth century, medical research was primarily based on empirical evidence and anecdotes. As the medicine has moved to become more evidence based, biostatistics has become ever more important in practicing medicine and for the education of tomorrow's doctors.<sup>1</sup> Biostatistical reasoning and modeling were of critical importance to the foundation

theories of modern biology. In the early 1900s, after the rediscovery of Mendel's work, the gaps in understanding between genetics and evolutionary Darwinism led to vigorous debate among biometricians. Statisticians then built models on statistical reasoning that helped to resolve these debates.<sup>2</sup> The role of biostatistics in medical education is now well recognized, and the curricula in almost all the medical

colleges and postgraduate institutions across globe has provision of teaching biostatistics<sup>3</sup> irrespective they are following traditional curriculum or integrated curriculum. The concept of integrated curriculum is nothing new. It's been around, in fact, since the 1800s and was advocated by well-known educational theorists (John Dewey and Meredith Smith). It has gained recent attention and most modern medical educationists think that it is the best way to teach.<sup>4-5</sup>

Biostatistics is a branch of applied statistics that is concerned with the application of statistical methods to medicine, clinical trials, demography, epidemiology, population estimation, modeling, community diagnosis and surveys.<sup>6</sup> Biostatistics is now considered as an essential tool in planning and delivery of health care systems. Integration is defined as a way to teach students how to break down barriers between subjects and make learning more meaningful & attractive. Integrated curriculum is an effective way to teach and learn because it corresponds with the way our brain works physiologically. According to M. Markus,<sup>7</sup> integrated curriculum is a great gift for experienced teachers. It's like getting a new pair of lenses that make teaching a lot more exciting and help us look forward into the next century.

Biostatistics is now being taught as a major subject in the medical curriculum. In a traditional curriculum, generally the course of biostatistics is spread over one academic year

study with formal final assessment. However, in an integrated curriculum it is generally taught as a longitudinal course spread over 4 – 6 months semester with either a formal final assessment or submitting a research project that depicts the knowledge of students. In this article, we will focus on “how biostatistics subject should be taught to undergraduate medical students in an integrated curriculum”.

*Biostatistics Course Induction Time in an Integrated Curriculum:* At undergraduate level,

there is a great difference in terms of organization of teaching and time allotted for biostatistics teaching in medical colleges globally. In United States, Canada, Australia and United Kingdom biostatistics is being taught in 1<sup>st</sup> and 2<sup>nd</sup> year of study.<sup>8</sup> In an integrated curriculum, the best time to start teaching biostatistics should be from 1<sup>st</sup> year of education, and biostatistical applications should be reinforced throughout during the basic sciences phase followed by a research project as course's final assessment. The core benefit of inducting biostatistics from 1st year level will strengthen the scientific reasoning of medical students and as they will reach the end of initial 2 ½ years of study their knowledge level will have increased unprecedentedly. Contra to it, if biostatistics is taught in one semester continuously it might bore the students because they are forced to grasp too much of knowledge in a limited time and will be tougher if it is followed by submitting a research project.

Secondly, with other parallel full time running courses the students might focus more on those subjects rather than learning biostatistics.

*Course of Biostatistics at Undergraduate Level in Integrated System:* In an integrated system biostatistics and research methods, subjects should be taught in-combination. Students should be imparted with the biostatistical and research methods knowledge throughout their first 2 ½ years of education. Learning the subject in-combination will not only help medical students to grasp necessary knowledge at undergraduate level but will also help them to develop research-oriented thinking at postgraduate level.

*Teaching Biostatistics in Integrated Curriculum:* In this section, some guidelines are given for teaching biostatistics in an integrated curriculum which may help to enhance the learning in this subject. The tutor must convince the medical students about the importance of learning biostatistics subject. As integration is a student-centered approach, one of the best ways of motivating the students is to expose them to the medical literature with examples of uses & abuses of biostatistics.<sup>9</sup>

In an integrated curriculum, the instructional method(s) should be problem-oriented instead of technique-oriented. The technique-oriented method is now old style which contains hand calculations, formulas and drawing critical regions etc.<sup>10</sup> Besides, problem-based approach

should be introduced which focuses more on interpretation. A survey conducted on biostatistics teaching showed that the students ostracized the subject because it was taught in the traditional way.<sup>11</sup>

Hand calculations should be avoided and for making the teaching interactive Computer based approach should be introduced. Nowadays a variety of biostatistical software's (SPSS, STATA, STATISTICA, NCSS and OPEN-EPI) are available. Introducing Computer Based Learning (CBL) will also enhance the student's motivation for learning. The use of computers in the teaching should be encouraged to allow the students to concentrate on the interpretation of the analysis.<sup>12</sup>

Reviewing literature is a pre-requisite for designing research. In an integrated curriculum, the tutor should educate the students about the methodology in searching literature and designing the research (research design, sampling technique / data collection and sample size etc). Moreover, the tutor should emphasize upon how results are interpreted and reported in biostatistical language.<sup>13</sup> Furthermore, the tutor should give special emphasis upon; how to write a research proposal and its scientific presentation. One of the main goals of teaching biostatistics is to analyze and interpret the data. In an integrated curriculum, it is the prime responsibility of the tutor to make the students understand about using accurate data analyzing technique(s).<sup>14</sup> The inappropriate use of



biostatistical methods & techniques may mislead the students understanding at a stage when their minds are fresh to grasp knowledge. The tutor should also strive to present a well-balanced combination of lectures, tutorials and practicals.<sup>15</sup>

## Conclusion

In an integrated curriculum biostatistics and research methods, course should be inducted in combination from 1st year of study. Curricula should be developed by content experts with taking care of the spiral approach. Students should be encouraged to conduct a research project of their own. Integrated curriculum is now considered the best way to teach so the tutors should get maximum benefit of teaching biostatistics under this system. The curriculum contents should be updated at the end of course every time based on students' feedback and recent advancements in the field. The computer based approach should be used to make the learning more interactive. Assessment must be aligned with teaching methods, curriculum outcomes and course objectives.

## References

1. Frank Emmert-Streib and Matthias Dehmer. *Medical Biostatistics for Complex Diseases*. Wiley-Blackwell 2010. ISBN 3-527-32585-9.
2. Terry Speed. *Microarray Gene Expression Data Analysis: A Beginner's Guide* 2003. Chapman & Hall/CRC.
3. Report of a Committee of the American Statistical Association Subsection on Teaching of Statistics in Health Sciences, Proposal for a core curriculum in medical biostatistics. *Clinical Pharmacology and Therapeutics* 1975;18:127-131.
4. Beane James A. *Curriculum Integration: Designing the Core of Democratic Education*. New York 1977: Teachers College Press.
5. Caine, Renate Nummela and Geoffrey Caine. *Making Connections: Teaching and the Human Brain* 1982.
6. Waqas Sami. *Biostatistics Education for Medical Students Pakistan's Perspective. Problems, Changing Approaches & Perceptions*. *Biomedica* 2010;26: 80-84.
7. M. Markus. *Media specialist*. Quoted in Shoemaker 1991, September, p. 797.
8. Jenny V Freeman, Steve Collier, David Staniforth and Kevin J Smith. *Innovations in curriculum design: A multi-disciplinary approach to teaching statistics to undergraduate medical students* *BMC Medical Education* 2008;8:28
9. Welch GE II, Gabbe SG. *Review of statistics usage in the American Journal of Obstetrics and Gynecology*. *Am J Obstet Gynecol* 1996;175(5):1138-41.
10. Boyle, C.R. "A Problem-Based Learning Approach to Teaching Biostatistics," *Journal of Statistics Education*, 1997;secure/v7n1/boyle.cfm.
11. Moussa, M. A. A. *Developments in the instruction of Biostatistics at the Kuwait University Health Science Centre in a decade*. *Teaching and Learning in Medicine*, 2002;14(3),194-198.
12. Tae Rim Lee. *Teaching Biostatistics to medical personnel with computer based supplement*. International Association for Statistical Education and International Statistical Institute. 2001, Printed in Granada, Spain.
13. Sahai, H. and Ojeda, M. M. *Teaching biostatistics to medical students and professionals: Problems and solutions*. *International Journal of Mathematical Education in Science and Technology*. 1999;30(2), 187-196.
14. Garfield J: *How students learn statistics*. *International Statistical Review* 1995;63:25-34.
15. Y.M. Lam. *Teaching Statistics to Medical Students in the New Medical School*. *Journal of the Hong Kong Medical Association*. 1986;38(4),196-198

**CASE STUDY**

## Chondroid Lipoma: A Rare Recently Described Benign Lipomatous Tumor

Salman T. Almalki<sup>\*1</sup>, Abdullah S. Alkhamiss<sup>2</sup>

\* For correspondence [bbbg126@hotmail.com](mailto:bbbg126@hotmail.com)

<sup>1,2</sup> Department of Pathology, King Fahad Medical City, Riyadh, Saudi Arabia

Submitted on: 18<sup>th</sup> March 2015, Accepted on: 6<sup>th</sup> September 2015

### Abstract

We present a very rare benign lipomatous tumor that has been described recently. A 51-year-old lady presented with a mass arising in her right thigh. This mass grew slowly during the several months ago. Radiologically, it was described as a well-circumscribed benign mass with scanty fatty component. Surgical excision and primary closure was performed for the patient. Histopathologic examination demonstrated a well-circumscribed mass formed by cords and nests of lipoblasts and lipocytes embedded in a prominent myxoid to hyalinized chondroid matrix. A panel of immunohistochemical (IHC) stains had been done. From the clinical, radiological, and pathological information, the diagnosis of chondroid lipoma was made. This case is considered a rare variant of lipoma and easily can be misdiagnosed as myxoid liposarcoma or extraskletal myxoid chondrosarcoma. So, the pathologist should pay attention when he/she comes across such cases.

### المخلص

نقدم حالة نادرة جدا لورم دهني حميد وهي من الحالات التي وصفت حديثا في المراجع. شكت سيدة تبلغ من العمر 51 سنة من ظهور ورم في فخذاها الأيمن وقد نما الورم ببطء خلال الشهور القليلة الماضية. اظهر التصوير الشعاعي وجود ورم حميد ذو حواف محددة جيدا مع وجود قليل من المكونات الدهنية، تم استئصال الورم. أجري الفحص المجهرى للنسيج المستأصل ووجد ان الورم ذو حواف محددة جيدا ويتكون داخليا من حبال واعشاش من الخلايا الدهنية ( lipoblasts and lipocytes) مغموسة في مصفوفة مخاطية غضروفية وزجاجية. كما تم عمل الفحص الصبغي مناعيا وكيميائيا. تم التشخيص على انه ورم دهني غضروفي. وتعتبر هذه الحالة نادرة من الأورام الدهنية وكان من الممكن بسهولة تشخيصها خطأ على انها حالة سرطان دهني مخاطي او سرطان غضروفي مخاطي. ولذلك علي طبيب علم الامراض وضع ذلك في الحسبان عند فحص مثل هذه الحالات.

### Introduction

Lipomatous tumors have a broad clinical and morphological spectrum ranging from benign cases to malignant ones. In the last few years, many variants of those lesions have been described. Chondroid lipoma is one of those variants. Although it is benign in nature, its histopathological morphology make it one of the great mimics of malignant soft tissue tumors.

### Case Report

51 years old female patient is presenting to the OPD with a right thigh swelling. This swelling

starts to appear several months ago. It is painless but slowly progressive. There is no decrease in weight or loss of appetite. On physical examination, there is a non-tender mass in the upper part of the right thigh. This mass is firm, deeply-seated, roughly oval, and not pulsated or fixed to the skin. CBC, LFT, RFT, and minerals are all within normal limits. The radiological examination shows well-defined mass contain only a small amount of fatty tissue distributed centrally & peripherally (Figure 1).

The mass have been excised and the gross picture shows a circumscribed lobulated tan-

yellowish mass measuring 5.2 x 4.3 x 3.8 cm (figure 2).




Figure 1: Radiological picture showing thigh with deep-seated well-circumscribed mass.




Figure 2: Gross picture showing a circumscribed tan-yellow lobulated mass.

Microscopically, the mass is well-circumscribed and formed by lobules separated by fibrous septa. Each lobule is formed by nests and cords of cells embedded in a chondroid-myxoid background Figure 3.

The cells have variable morphology: some of them are mature adipocytes with single large intracytoplasmic vacuole, others have multivacuolated cytoplasm resembling lipoblasts Figure 4 and 5.




Figure 3: Low-power view demonstrating lobular pattern with a myxoid background.




Figure 4: Intermediat-power view showing mature fat cells with degenerative changes and the surrounding chondromyxoid matrix.




Figure-5: High-power veiow showing mature fat cells and lipoblasts embedded in a chondroid matrix. Some lacunae-like cells are also present.



Some of the cells have a pericytoplasmic clear zone (lacunae-like), making it resembling chondrocytes Figure 5. Mild nuclear pleomorphism and rare scattered mitoses are noted Figure 6.




Figure 6: High-power view showing many lipoblasts with mild nuclear pleomorphism embedded in a chondroid matrix. (H&E x400).




Figure-7: The cells are diffusely positive for vimentin (immunohistochemistry stain x200).

No chicken-wire blood vessels seen in the background. IHC stains shows that the tumor cells are positive for vimentin (figure 7) & S100 (Figure 8) and negative for CK, EMA, CD34, SMA, desmin, & HMB45. Based on the previous findings, a diagnosis of chondroid

lipoma is made. The patient had been discharged with a follow-up appointment.




Figure 8: The cells are focally positive for S100 (immunohistochemistry stain x200).

### Discussion

Lipomatous tumors are one of the most common soft tissue tumors that usually diagnosed by the pathologists. They have a wide range of morphological and behavioral spectrum from benign things (lipoma) to malignant one (liposarcoma).<sup>1-2</sup> Chondroid lipoma is a very rare variant of lipomatous tumor.<sup>3</sup> It had been firstly described by Meis and Enzinger<sup>4</sup> in 1993. They describe it as a lipoma contains two histological components: a) adipose tissue (containing mature fat cells & lipoblasts), and b) cartilaginous tissue (containing chondroblastes & hyaline matrix).<sup>4</sup> Now, according to the World Health Organization (WHO) 2002,<sup>3</sup> chondroid lipoma is defined as "a unique & recently recognized benign adipose tissue tumor containing lipoblasts, mature fat cells, and chondroid matrix". Why this tumor occurs? still unknown, but there are two

hypothesis regarding to the origin of the cartilaginous component of this tumor: a) it represents a metaplastic changes in an ordinary lipoma due to a mechanical stress, and b) it arising from the multipotential stem cells under the influence of some growth factors that are present in this lesion.<sup>3,5</sup> Regarding to the age group, usually this type of tumor will affect adult people in their third and fourth decades<sup>3,5</sup> but it can arise in pediatric age group also.<sup>7</sup> It arise more in females.<sup>3,6</sup> Proximal extremities and limb girdles are the most common affected sites, however, trunk and head & neck areas can be affected too.<sup>3</sup> It presented as a painless, slowly growing, superficial or deeply seated soft tissue mass.<sup>3,6</sup>

Grossly, it is a well-circumscribed lobulated tan-yellow mass with a size range of 2-7 cm.<sup>3,6</sup> Microscopically, the tumor is a well-circumscribed lobulated mass with lobules separated by fibrous septa.<sup>3,6</sup> Its histological hallmarks are nests and cords of abundant uni- and multivacuolated lipoblasts embedded in a prominent myxoid to hyalinized chondroid matrix admixed with a variable amount of mature adipose tissue.<sup>3</sup> Lipocytes are also present. NO/mild pleomorphism and rare mitosis are accepted. Some small cells with lacunae-like zone are present. No chicken-wire blood vessels in the background but some scattered thick wall blood vessels are noticed.<sup>3,6</sup> The cells contain both glycogen and lipid which highlighted by PAS and oil-red O stains,

respectively. Reticulin stain shows that there is a pericellular reticulin fiber. Immunohistochemically, the cells are positive for S100, vimentin, CD68, and rarely for CK. They are negative for EMA, HMB45, SMA, MSA, GFAP, and CD57<sup>1,3,6</sup>. Ki67 proliferation index is less than 1%. Cytogenetically, some of chondroid lipomas have t (11;16) which resulting in C11 or f95-MKL2 fusion oncogene.<sup>2-3,6</sup> How this fusion will lead to the formation of this tumor? Still no clear answer.

The differential diagnosis of chondroid lipoma includes benign and malignant tumors.<sup>1-3,6,8-9</sup> Myxoid liposarcoma is one of the most mimickers for chondroid lipoma but it can be differentiated by its infiltrative pattern, the presence of chicken-wire blood vessels, and the presence of t (12;16) or t (12;22) which will lead to the fusion of DDIT3 with FUS or EWS1 genes, respectively.<sup>2-3,6</sup>

Myxoid chondrosarcoma is another mimickers for chondroid lipoma but it can be differentiated by its infiltrative pattern, absence of the cytoplasmic vacuoles, absence of the fatty components, and the presence of chromosomal translocation that causing fusion of NR4A3 in chromosome 9. Spindle cell lipoma is one of the differential diagnosis of chondroid lipoma but it have ropey collagen fibers, lack of chondroid background, has diffusely expressed CD34 in the spindle cell component, and a deletion involving chromosomes 13q and 16q.<sup>2-3,6</sup>



It is important to know this type of benign tumor to avoid misdiagnosing it as a sarcoma and the subsequent overtreatment of the patients with radical surgery, chemotherapy, and radiotherapy.<sup>1-4,6,8,10-11</sup> Chondroid lipoma will be cured with simple excision<sup>3</sup>. It will not recur or metastasis.<sup>3</sup>

## Conclusion

Chondroid lipoma is a rare variant of lipoma with histomorphological features that mimic myxoid liposarcoma and extraskeletal myxoid chondrosarcoma. Attention should be paid from the pathologist when he/she evaluates such cases, particularly on core biopsy sample.

## References

1. Gokhale U, Pillai GR, Varghese PV, Samarsinghe D. Chondroid lipoma: a case report. *Oman Medical Journal* 2008; 23(2),65-69
2. De Vreeze R, Van Coevorden F, Boerrigter L, Nederlof PM, Haas RL, Bras J, et al. Research article delineation of chondroid lipoma: an immunohistochemical and molecular biological analysis. *Hindawi Publishing Corporation Sarcoma* 2011; Article ID 638403:5.
3. Fletcher C.D.M., Unni K.K., Mertens F. (Eds.): *World Health Organization (WHO) Classification of Tumours. Pathology and Genetics of Tumours of Soft Tissue and Bone*. IARC Press: Lyon 2002.
4. Meis JM, Enzinger FM. Chondroid lipoma. A unique tumor simulating liposarcoma and myxoid chondrosarcoma. *Am J Surg Pathol* 1993;17(11):1103-1112.
5. Hwang HS, Lee WJ, Lim HK, Chun HK, Ahn GH. Chondrolipoma in the pelvic cavity: a case report. *Korean J Radiol* 2008;9:563-567.
6. Thway K, Flora RS, Fisher C. Review article chondroid lipoma: an update and review. *Annals of Diagnostic Pathology* 2012;16:230-234.
7. Kindblom LG. Lipomatous tumors-how we have reached our present views, what controversies remain and why we still face diagnostic problems: a tribute to Dr Franz Enzinger. *Adv Anat Pathol* 2006;13(6):279-285.
8. Gomez-Ortega JM, Rodilla IG, De Lerma JM. Chondroid lipoma a newly described lesion that may be mistaken for malignancy. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology* 1996;18(5):586-589.
9. Lakshmiah SR, Scott KW, Whear NM, Monaghan A. Chondroid lipoma: a rare but diagnostically important lesion. *International Journal of Oral and Maxillofacial Surgery* 2000; 29(6):445-446.
10. Nielsen GP, O'Connell JX, Dickersin GR, Rosenberg AE. Chondroid lipoma, a tumor of white fat cells. A brief report of two cases with ultrastructural analysis. *Am J Surg Pathol* 1995;19(11):1272-1276.
11. Mentzel T, Remmler K, Katenkamp D. Chondroid lipoma. Clinicopathological, immunohistochemical, and ultrastructural analysis of six cases of a distinct entity in the spectrum of lipomas. *Pathologie* 1999; 20(6): 330-334.

**MEDICAL NEWS**

# World Health Organization

## WHO Health Topic: Are we practice a safe health care?

All health care providers are dedicating themselves to provide a good health care to their patient with no harm, but remains the fact that patients are harmed daily in the course of receiving health care all over the world. Patient safety is a serious global public health issue. Estimates show that in developed countries as many as 1 in 10 patients is harmed while receiving care.

Patient safety has received increased attention in recent years, but mostly with a focus on the epidemiology of errors and adverse events, rather than on practices that reduce such events. Therefore, we need to address this fact and do our best to correct the problems that leads to unsafe practice, as all patients have a right to effective, safe care at all times. WHO considered patient safety as a fundamental principle of health care. Every point in the process of care-giving contains a certain degree of inherent unsafety. Adverse events may result from problems in practice, products, procedures or systems. Patient safety improvements demand a complex system-wide effort, involving a wide range of actions in performance improvement, environmental safety and risk management,

including infection control, safe use of medicines, equipment safety, safe clinical practice and safe environment of care.

Recently, the solutions for insuring patient safety offer a more constructive approach; one important solution is the work of the caregivers together as a team, how effectively they communicate with one another and with patients, and how carefully the care delivery processes and supporting systems of care are designed. With the growing recognition of safety problems in health care, it is now time to create and disseminate “Solutions” for patient safety. WHO stated some guidelines on patient safety in some health practices, it is under the following headings:

- Best practices for injections and related procedures toolkit
- Blood donor selection
- Natural ventilation for infection control in health-care settings
- Screening donated blood for transfusion-transmissible infections: recommendations
- Towards 100% voluntary blood donation

- WHO guidelines on hand hygiene in health care

Not only that but also WHO published a Patient Safety Curriculum Guide under the title “The Multi-professional Patient Safety Curriculum Guide” It was released in October 2011 and it promotes the need for patient safety education. The comprehensive guide assists universities and schools in the fields of dentistry, medicine, midwifery, nursing and pharmacy to teach patient safety. It also supports the training of all health-care professionals on priority patient safety concepts and practices.

The World Health Organization launched the World Alliance for Patient Safety and identified six action areas in 2005. One of these action areas is the development of “Solutions for Patient Safety”. In the same year, the Joint Commission and Joint Commission International were designated as a WHO Collaborating Centre for Patient Safety Solutions, to initiate and coordinate the work of developing and disseminating solutions for patient safety. The output from this component of the World Alliance will be delivered to the global health-care community as “Patient Safety Solutions”. Within the foregoing context, the term “Patient Safety Solution” is defined as: Any system design or intervention that has demonstrated the ability to prevent or mitigate patient harm stemming from the processes of health care.

Inaugural Patient Safety Solutions:

1. Look-Alike, Sound-Alike Medication Names
2. Patient Identification
3. Communication During Patient Hand-Overs
4. Performance of Correct Procedure at Correct body site
5. Control of Concentrated Electrolyte Solutions

6. Assuring Medication Accuracy at Transitions in Care
7. Avoiding Catheter and Tubing Mis-Connections
8. Single Use of Injection Devices
9. Improved Hand Hygiene to Prevent Health Care-Associated Infection

International patient safety goals:

- Identify patient correctly
- Improve effective communication
- Improve the safety of high alert medication
- Eliminate wrong site, wrong patient and wrong procedure surgery
- Reduce the risk of health care acquired infection
- Reduce the risk of patient harm resulting from fall

Fortunately, authorities in many countries are reforming health care in terms of higher quality and the elimination or correction of practices that are known to be unsafe or wasteful. Also let us not forget the role of patients and their families in improving the patient safety, they are becoming increasingly skilled in accessing information to make personal health care decisions about treatments and their choice of providers, and demanding safer care as well. Health-care practitioners are also becoming more proficient at incorporating evidence-based knowledge into their clinical decision-making practices. One of the role of patients and their families in improving the safety is the patient advocates use past experiences to change the future, and here is an example of a lady who lost her son and she dedicated herself to advocate patient safety. When visiting a doctor, patients expect a listening ear, an accurate diagnosis and timely treatment. This woman encountered the opposite, a flawed health system that lacked the capacity to respond to her 21

years old son deteriorating health. His personality changed. He was moody and constantly tired. He consulted a doctor for persistent bone pain. Tests revealed high calcium levels and a plasma creatinine level indicating kidney failure. The results required immediate referral, which did not happen. Symptoms continued for another 2 years without action. On his last visit to his doctor, tests showed critically high levels of calcium in his blood, levels later described by peer reviewers as "inconsistent with life". The results were communicated by telephone and written on a post-it note by a nurse, who believed that the patient had hypercalcaemia. Ignoring the nurse's diagnosis, the doctor omitted any reference to calcium and only included results in his referral letter that supported the incorrect diagnosis. The post-it note was stuck to the back of the referral letter and was not seen by the hospital until 6 weeks after the patient death. Unaware of his high calcium levels, the patient was diagnosed with nephritis and transferred to another hospital for treatment. Because it was a weekend, on-call senior personnel were not advised of the patient's condition and an aggressive treatment plan was not started. Hours later, the patient suffered a heart attack and could not be resuscitated. His cause of death was primary hyperparathyroidism, multi-organ failure and hypercalcaemia. Calcium was being channelled into the patient's blood stream instead of his bones, putting strain on his heart. A simple surgery, with a 96% success rate, would have saved his life.

"Adverse events happen to real people," the mother says. "my son was more than a statistic; he was more than a medical condition. He was a real person, a young man, full of life. But above all, he was my beautiful boy."

In 2004, WHO launched a patient safety programme to develop policies and practices aimed at preventing harm to patients worldwide. Because the patient voice is essential to advocating for change, WHO also formed Patients for Patient Safety (PFPS) – a patient-lead, global network of patients and patient organizations who advocate for patient safety.

"Because patients and caregivers see things that busy health-care workers do not, it's essential for us to include them at the center of all the work we do."

"The patient is the single entity which is present throughout the full continuum of care," stresses Margaret. "A variety of professionals dip in and out of the care pathway, but the patient is the one constant and a useful resource in relation to diagnosis and treatment. Importantly, the patient is the individual with the greatest vested interest in the outcome."

## MEDICAL QUIZ

**Moattar R Rizvi, Assistant Professor, Physiology**

An 18 years old female patient presents to the emergency department stating that over the past 24 hours, she has developed left sided facial weakness and drooping. Her mouth turns towards the side if she smiles and she is unable to close her left eye.


She also complains that she is unable to move both upper and lower side of her face. She stated that she has had a herpes simplex (viral) infection.

O/E of Vital signs –

Vital Measurements	Recorded values	Normal values
Blood Pressure (mm of Hg)	124/80	120/80–140/90
Temperature (°F)	98.1	97 - 99
Respiratory Rate (per min)	15	12 - 20
Pulse (per min)	78	60 - 100

Systemic examination was unremarkable and investigations like skull xray, CT and MRI were normal.

The physician diagnosed the case as Bell's palsy and prescribed an antiviral medication along with a short course of corticosteroids.



**Depending on this case, please answer the below mentioned questions.**

- Bell's palsy affects one of the following cranial nerves :
  - Trigeminal
  - Facial
  - Glossopharyngeal
  - Hypoglossal
- This patient will complain of suppressed gustatory sensation from which of the following region:
  - Anterior two thirds of the tongue
  - Posterior one third of the tongue
  - Epiglottis and pharynx
  - All of the above
- The muscles of mastication (chewing) are ..... in Bell's palsy.
  - Affected
  - Unaffected (spared)
  - Partially affected
- Tactile sensation from the face will be unaffected in Bell's palsy.
  - True
  - False
- The patient has complained of dry eyes (left eye). Which of the following options given below is the correct explanation?
  - Impaired innervations of the lacrimal gland
  - Inability to close the eyes leading to evaporation of tears
  - Both of the above
  - None of the above

Tick the correct statement:

- Bell's palsy is due to lower motor neurone lesion. T    F
- Bell's palsy is idiopathic in nature. T    F
- The benefits of acyclovir is definitive on the patient T    F
- About 50% of people get better with no treatment T    F
- Gustatory sweating is a recognized complication T    F



**UPCOMING CONFERENCES**

	<b>Conference List</b>	<b>Date/Venue</b>
<b>Septembers, 2015</b>	<b>3rd International Oncology Conference</b>	<b>10 – 11 September, 2015, Abu Dhabi, UAE</b>
	<b>The 8th Pan Arab Radiology Conference (ARC 8) The 10<sup>th</sup> Jordanian Radiology Society Conference</b>	<b>10 - 13 September, 2015, Amman, Jordan</b>
<b>October, 2015</b>	<b>4th International Conference on Surgery</b>	<b>5 – 7 October, 2015, Dubai, UAE</b>
	<b>Mental Health Congress (WFMH 2015)</b>	<b>16 – 19 October, 2015, Cairo, Egypt</b>
	<b>3rd Annual Pediatric Surgical Innovation Symposium</b>	<b>23 October 2015, Washington, D.C., USA</b>
	<b>4th International Conference on Orthopedics &amp; Rheumatology</b>	<b>October 26-28, 2015, Baltimore, USA</b>
<b>November, 2015</b>	<b>International Paediatric Medical Congress</b>	<b>12 – 14 November, 2015, Dubai, UAE</b>
	<b>The IRES – 31st International Conferences on Medical and Health Science (ICMHS)</b>	<b>16th to 16th November 2015 Riyadh, Saudi Arabia</b>
	<b>Abu Dhabi International Conference in Dermatology &amp; Aesthetics</b>	<b>19 – 20 November, 2015, Abu Dhabi, UAE</b>
	<b>XXI. World World Congress of Echocardiography and Cardiology</b>	<b>20-22 November, 2015, Istanbul, Turkey</b>
	<b>1st Saudi Epidemiology Conference</b>	<b>November 24-26, 2015 Hotel Park Hyatt – Jeddah</b>

	<b>Global Summit and Medicare Expo on Head &amp; Neck Surgery</b>	<b>November 30-December 01, 2015, Atlanta, USA</b>
<b>December, 2015</b>	<b>4th Saudi Heart Failure Conference</b>	<b>December 4, 2015 Jeddah, Saudi Arabia</b>
	<b>International Conference and Exhibition on Cosmetic Dermatology and Hair Care</b>	<b>December 7-8 2015, Philadelphia, USA</b>
	<b>3rd International Endoscopy Symposium</b>	<b>09-10 Dec 2015, Riyadh, KSA</b>
<b>January, 2016</b>	<b>International Conference On Infectious Disease</b>	<b>26-27 Jan 2016, Jeddah, KSA</b>
	<b>18th International Conference on Diet, Gut Microbiology and Human Health</b>	<b>January 26 - 27, 2016 Jeddah, KSA</b>
<b>February</b>	<b>12th International Conference on Medical and Biosciences (ICMBS)</b>	<b>22nd February 2016 Riyadh, Saudi Arabia</b>
<b>March, 2016</b>	<b>5th World Congress on Neurology and Therapeutics</b>	<b>March 14-16, 2016, London, UK</b>
	<b>7th World Cardiothoracic Meeting</b>	<b>March 29-30, 2016, Atlanta, USA</b>
	<b>Annual Pediatric Emergency Medicine Conference</b>	<b>March 29-31, 2016, Atlanta, USA</b>
	<b>4th International Conference on Pediatrics</b>	<b>March 29-31, 2016, Atlanta, USA</b>
<b>May</b>	<b>Saudi Health 2016</b>	<b>16 - 18 May 2016, Riyadh, KSA</b>

# GUIDELINES FOR MANUSCRIPT PREPARATION

## **A. TYPES OF MANUSCRIPTS**

### **I. ORIGINAL MANUSCRIPTS**

Manuscripts submitted in this category are expected to be concise, well organized, and clearly written. The maximum length is 5000 words, including the abstract, references, tables, and figure legends. The maximum length is 5000 words, including the abstract, references, tables, and figure legends.

- The structured abstract must not exceed 250 words.
- The title must not exceed 130 characters.
- A maximum of 4 tables and 4 figures is allowed.
- References should not exceed a maximum of 100.
- The abstract must be organized as follows:
  - Background & Aims
  - Methods
  - Results
  - Conclusions
- Do not use abbreviations, footnotes or references in the abstract.
- An electronic word count of the abstract must be included.
- Three to ten key words at the end of the abstract must be provided.

The manuscript must be arranged as follows:

- Title page
- Abstract
- Introduction
- Materials and methods (or Patients and methods)
- Results
- Discussion
- Acknowledgements
- References
- Tables
- Figure legends
- Figures

Acceptance of original manuscripts will be based upon originality and importance of the investigation. These manuscripts are reviewed by the Editors and, in the majority of cases, by two experts in the field. Manuscripts requiring extensive revision will be at a disadvantage for publication and will be rejected. Authors shall be responsible for the quality of language and style and are strongly advised against submitting a manuscript which is not written in grammatically correct English. The Editors reserve the right to reject poorly written manuscripts even if their scientific content is qualitatively suitable for publication. Manuscripts are submitted with the understanding that they are original contributions and do not contain data that have been published elsewhere or are under consideration by another journal.

### **II. REVIEW ARTICLES**

Review articles on selected clinical and basic topics of interest for the readers of the Majmaah Journal of Health Science will be solicited by the Editors. Review articles are expected to be clear, concise and updated.

- The maximum length is 5000 words, excluding the summary, references, tables, and figures.
- References should not exceed a maximum of 150.
- The inclusion of a maximum of 4 high-quality tables and 4 colored figures to summarize critical points is highly desirable.
- Review articles must be accompanied by a title page and a summary.

- Reviews should include at least one Key Point Box, with a maximum of 5 bullet points, that briefly summarizes the content of the review.

Review articles are reviewed by the Editors and may be sent to outside expert reviewers before a final decision for publication is made. Revisions may be required.

### III. EDITORIALS

This section consists of invited brief editorial comments on articles published in the Majmaah Journal of Health Science

The length of an editorial should not exceed 1500 words, excluding references.

- A maximum of 1 table or 1 figure is allowed.
- References should not exceed a maximum of 20.
- A title page must be provided.

### IV. CASE REPORTS

Case reports would be only accepted if they represent an outstanding contribution to the Etiology, pathogenesis or treatment of a specific condition.

- The maximum length is 3000 words, including the summary and references.
- A maximum of 2 tables and 2 figures is allowed.
- References should not exceed a maximum of 15.
- A title page must be provided.

### V. LETTERS TO THE EDITOR

Letters to the Editor will be considered for publication if they are related to articles published in recent issues of Majmaah Journal of Health Science. Occasionally, Letters to the Editor that refer to articles not published in Majmaah Journal of Health Science will be considered.

The length of a Letter to the Editor should not exceed 800 words.

- A maximum of 1 table or 1 figure is allowed.
- References should not exceed a maximum of 10.
- No more than 4 Authors may appear in the author list.

### VI. COMMENTARIES

International commentaries will be solicited by the Editors only.

- Commentary articles should not exceed a maximum of 800 words, excluding tables or figures.
- A maximum of 1 table or 1 figure is allowed.
- References should not exceed a maximum of 10.
- A title page must be provided.

## B. MANUSCRIPT SUBMISSION

### ORGANIZATION OF THE MANUSCRIPT

- The submitted manuscript must be typed double-spaced throughout and numbered (including references, tables and figure legends). Preferably using a "standard" font (we prefer Times/Arial 12).
- For mathematical symbols, Greek letters, and other special characters, use normal text. The references must be in accordance with the Vancouver reference style (see References).
- Approved nomenclature for gene and protein names and symbols should be used, including appropriate use of italics (all gene symbols and loci, should be in italics) and capitalization as it applies for each organism's standard nomenclature format, in text, tables, and figures.
- Full gene names are generally not in italics and Greek symbols are not used. Proteins should not be italicized.
- Improperly prepared manuscripts will not be entered into the peer review process and will be sent back to the author for correction.

### TITLE PAGE MUST CONTAIN:

- A title of no more than 130 characters.



- Running title (not to exceed 60 characters)
- Names of the Authors as it should be published (first name, middle initial, last name)
- Affiliations of all authors and their institutions, departments, or organizations (use the following symbols in this order to designate authors' affiliations: \*, †, ‡, §, ¶, ||, #, \*\*, ††, ‡‡, §§, ¶¶, || ||, ##).
- Name, address, telephone and fax numbers, and electronic mail address of the corresponding Author.
- Electronic word count.
- Number of figures and tables.
- List of abbreviations in the order of appearance.
- Conflict of interest.
- Financial support.

**Animal trials:** Manuscripts reporting experiments using animals must include a statement giving assurance that all animals received human care and that study protocols comply with the institution's guidelines. Statistical methods used should be outlined.

**Human trials:** Manuscripts reporting data from research conducted on humans must include a statement of assurance in the methods section of the manuscript reading that:

1. Informed consent was obtained from each patient included in the study and
2. The study protocol conforms to the ethical guidelines of the 1975 declaration of helsinki as reflected in a priori approval by the institution's human research committee.

**Randomized controlled trials:** Any paper that is a randomized control trial should adhere to the guidelines that can be found at the following web-site: [www.consort-statement.org](http://www.consort-statement.org). The checklist should be printed out and faxed to the Editorial office at the time of submission. The trial registration number must be included on the title page of the manuscript reporting a registered clinical trial. Failure to do so will prevent entry to the peer review process.

**Drugs and chemicals:** Drugs and chemicals should be used by generic name. If trademarks are mentioned, the manufacturer's name and city should be given. All funding sources supporting the work, either public or private, especially those from pharmaceutical companies, must be provided.

**Genetic Sequence data:** In papers reporting a novel DNA or amino sequence, verification that the data have been or will be submitted either to Gen-Bank or EMBL is required. Please provide this verification and the accession number in the covering letter.

## REFERENCES

References must be in accordance with the Journal of Hepatology reference style. References are ordered as they appear in the text and citation numbers for references are placed between "brackets" ("[ ]") in the text as well as in the reference list.

Authors should be listed surname first, followed by the initials of given names (e.g. Bolognesi M). If there are more than six authors, the names of the first six authors followed by et al. should appear.

Titles of all cited articles are required. Titles of articles cited in reference list should be in upright, not italic text; the first word of the title is capitalized, the title written exactly as it appears in the work cited, ending with a full stop. Journal titles are abbreviated according to common usage, followed by Journal years, semicolon (;) before volume and colon (:) before full page range (see examples below).

All articles in the list of references should be cited in the text and, conversely, all references cited in the text must be included in the list.

Personal communications and unpublished data should be cited directly in the text by the first Author, without being numbered. Please make sure you have the latest, updated version of your reference management software to make sure you have the correct reference format for Majmaah Journal of Health Science.

*An example of how references should look within the text:*



"HVPG was measured by hepatic vein catheterization using a balloon catheter according to a procedure described elsewhere [14, 15] and used as an index of portal hypertension [16]."

*An example of how the reference list should look:*

[14] Merkel C, Bolognesi M, Bellon S, Zuin R, Noventa F, Finucci G, et al. Prognostic usefulness of hepatic vein catheterization in patients with cirrhosis and esophageal varices. *Gastroenterology* 1992;102:973-979.

[15] Groszmann RJ, Wongcharatrawee S. The hepatic venous pressure gradient: anything worth doing should be done right. *Hepatology* 2004;39:280-282.

## FIGURES

A maximum of 4 figures is allowed

(This can be modified if needed by Editorial board).

- Figures will be often, but not always, re-designed by graphic designers. By signing and transferring the Copyright Agreement to MJHS, the author gives permission to the graphic designers to alter the visual aspect of any figures, tables, or graphs. The scientific content of figures will not be altered. Please provide this information with your covering letter.
- All graphics submitted to Majmaah Journal of Health Science should be sent at their actual size, which is 100% of their print dimension and in portrait orientation.
- Two standard widths are used and figures should fit in one (8.5 x 23.5 cm) or two (17.5 x 23.5 cm) columns
- Figures should be supplied in the following preferred file formats: PDF (\*.pdf), Power Point (\*.ppt), Adobe Illustrator (\*.ai, \*.eps), Photoshop (\*.psd) files in grayscales or in RGB color mode. It is highly recommended that figures not be sent in JPG (\*.jpg) format.
- Photographs (scans, immunofluorescences, EM, and histology images) should be submitted as: 1. TIFF (\*.tif) with a resolution of at least 300 pixels per inch, or
- Illustrator compatible EPS files with RGB color management (\*.eps),
- Photoshop (\*.psd) or PDF (\*.pdf) files (grayscales or RGB) at the appropriate resolution, which is:
  1. 300 dpi for color figures
  2. 600 dpi for black and white figures
  3. 1200 dpi for line-art figures
- For all photomicrographs, where possible, a scale should appear on the photograph. Photographs of identifiable patients should be accompanied by written permission to publish from patient(s).
- Furthermore, panel lettering should be in Arial bold 14 pt, capitalized and no full stop (A, B) while lettering in figures (axes, conditions), should be in Arial 8 pt, lower case type with the first letter capitalized and no full stop. No type should be smaller than 6 pt.

## TABLES

A maximum of 4 tables is allowed

(This can be modified if needed by Editorial board)

- Tables should be provided as Word files (\*.doc) or Illustrator/InDesign (\*.ai, \*.eps, \*.indd) compatible files. No TIFF and JPG files are acceptable for table submission.
- When submitting tables in Microsoft Word table function, no tab, space or colors should be used. Tables should contain a maximum of 10 columns.
- Tables submitted in landscape orientation will not be accepted. Tables should include a title, table legend, and if necessary footnotes.
- Include tables in the submitted manuscript as a separate section.

## FIGURE LEGENDS

- Figure legends should be listed one after the other, as part of the text document, separate from the figure files.
- Please do not write a legend below each figure. Each figure legend should have a brief title that describes the entire figure without citing specific panels, followed by a description of each panel, and the symbols used.
- Enough information should be provided in the figure legend text to permit interpretation of figures without reference to the text; but should not contain any details of methods, or exceed 100 words.
- The abbreviated word for figure "Fig." should be typed and bolded, followed by the figure number and a period

(i.e. "Fig. 1."). Every figure legend should have a Title written in bold.

- If a figure contains multiple sections (i.e. A, B, C, D) the letter for these subsections should be in capital letters. Within the figure legend text the capital letters should be surrounded by parenthesis [i.e. (A)(B)(C)(D)].
- Figures should be numbered according to the order of citation.

**Supplementary material:** Supplementary material, not for review, is acceptable. Supplementary material can be submitted as (\*.mov), (\*.avi), (\*.mpeg), or (\*.gif) files. Please note that the size limit for these items is 10 MB per file.

## ENGLISH

Authors may be asked to contact professionals regarding the correction of the English content of manuscripts either before or after acceptance. This expense will be the responsibility of the Authors.

## C. REVIEW PROCESS

Authors should be aware that manuscripts will be screened upon submission. Only the manuscripts which fully comply with the submission requirements outlined and in which the level of English is of an acceptable standard will enter the peer review process.

### First submission

Once successful submission of a manuscript has taken place, an acknowledgement will be sent by e-mail to the Corresponding Author on the manuscript. All subsequent correspondence will be with the designated Corresponding Author. The number of the manuscript should be used by the Authors in all communications with the Editorial Office. All the manuscripts will be reviewed by the Editors and, in some cases, by other expert reviewers. After review, the corresponding Author will be notified by letter of the decision taken by the Editor(s). This letter will be accompanied in most, but not all, cases by the comments of the reviewers. This letter will be sent via e-mail.

### Resubmission of manuscripts

In some cases, Authors will be invited to submit a revised version of the manuscript for further review. This invitation does not imply, in any case, that the revised version will be accepted for publication. In general, revised manuscripts must be received in the Editorial Office within four months of the date of the first decision. Authors should submit the resubmitted manuscript with all changes underlined. The resubmitted manuscript should be accompanied by a cover letter stating that the manuscript has been revised according to the comments made by the Editor and the Reviewers. Figures and tables must be uploaded. Please ensure that a separate point by point response to the reviewers is included with the covering letter. Please do not send revised manuscripts to the Editorial Office via e-mail. Revised manuscripts should be mailed to site of Majmaah Journal of Health Sciences at [mjhs@mu.edu.sa](mailto:mjhs@mu.edu.sa)

## PROOFS

Proofs will be made available to the author(s) to be checked. It is the responsibility of the author(s) to make sure that the quality and accuracy of the manuscript, figures, and tables in the proofs is correct. Authors should return their proofs within 48 hours, by fax or e-mail if the corrections are minor, to expedite publication. Further changes or additions to the edited manuscript after these corrections cannot be accepted.

## COVER ILLUSTRATIONS

Cover illustrations will be chosen by the Editors. Authors are highly encouraged to submit high quality color figures and images suitable for publication on the cover at the time of submission of the manuscript.

## REPRINTS


Reprints must be ordered in advance. An order form indicating the cost of the reprints is sent from the Publisher with page proofs. Reprint orders, payments, and inquiries must be forwarded to the Publisher, not to the Editorial Office.

## ADVERTISEMENTS

Information about advertisements in Majmaah Journal of Health Science can be obtained from the Publisher.

Copyright assignments, financial disclosures, and Institutional Review Board/Animal Care Committee Approval. Upon article acceptance, the corresponding author will be contacted and asked to submit the above forms. It is the author's responsibility to make sure these forms are signed and duly returned to the editorial office via fax. If these forms are





not received the manuscript will NOT be published.

#### **Drug Declaration/Conflict of Interest Form**

This form should be printed out and the suitable statement chosen among the listed ones (A-G). It should then be signed by the corresponding author and faxed to the Editorial Office at +41 22 510 24 00. If this form is not received the paper will NOT be published.

### ***Methodological & Statistical instructions for Authors submitting manuscripts to the Majmaah Journal of Health Science***

The manuscripts should include a complete and detailed description of what was done. This includes a description of the design, measurement and collection of data, the study objective and major hypotheses, type and source of subjects, inclusion and exclusion criteria and measures of outcome, number of subjects studied and why this number was chosen. Any deviation from the study protocol should be stated. The baseline characteristics of any compared groups should be described in detail and -if necessary -adjusted for in the analysis of the outcome.

For randomized clinical trials the following should also be clearly documented: treatments, sample size estimation, method of random allocation and measures taken for maintaining its concealment including blinding, numbers treated, followed-up, being withdrawn, dropping out, and having side effects (numbers and type). The statistical methods used should be relevant and clearly stated. Special or complex statistical methods should be explained and referenced.

Complex analyses should be performed with the assistance of a qualified statistician. Unqualified use of such analyses is strongly discouraged. The underlying assumptions of the statistical methods used should be tested to ensure that the assumptions are fulfilled.

For small data sets and if variable distributions are non-normal, distribution free (non-parametric) statistical methods should be used. The actual p values - whether significant or not - should always be presented (not NS). Confidence intervals convey more information than p values and should be presented whenever possible. Continuous variables can always be summarized using the median and range which are therefore preferred. Only in the infrequent case of a Normal distribution are the mean and standard deviation (SD) useful. Complex analyses (including Cox and logistic regression analysis) should be presented in sufficient detail: i.e. variable scoring, regression coefficients, standard errors and any constants. Odds-ratios or relative risks are not sufficient documentation of such analyses. The handling of any missing values in the data should be clearly specified. The number of statistical tests performed should be kept at a minimum to reduce spurious positive results. Explorative (hypothesis generating) analyses without confirmation using independent data are discouraged. Figures showing individual observations e.g. scatter plots are encouraged. Histograms may also be useful. Tables should indicate the number of observations on which each result is being based





# MJHS

مجلة الجامعة للعلوم الصحية

MAJMAAH JOURNAL OF HEALTH SCIENCES  
A JOURNAL PUBLISHED BY MAJMAAH UNIVERSITY

## Editor in Chief

Prof. Mohammad Othman Al-Rukban

## MJHS EDITORIAL ASSISTANTS

Dr. Khalid Tohami

Dr. Fuzail Ahmad

Dr. Moattar Raza Rizvi

Mr. Waqas Sami

## MJHS Office

MAJMAAH JOURNAL OF HEALTH SCIENCES  
Basic & Health Sciences Research Centre  
Deanship of Research, Majmaah University  
Ministry of Education,  
Kingdom of Saudi Arabia  
Post Box 66, Al Majmaah 11952  
E-Mail: [mjhs@mu.edu.sa](mailto:mjhs@mu.edu.sa)

Website: <http://mu.edu.sa/en/departments/Majmaah-journal>

## INVITATION TO PUBLISH

Dear Authors,

Majmaah Journal of Health Sciences is accepting original articles, review articles and letters to editors for publishing in forthcoming issue. Kindly go through the guidelines for submission and submit your manuscripts online at [mjhs@mu.edu.sa](mailto:mjhs@mu.edu.sa)



Copyrights© 2015 by Majmaah University. All right reserved.

Reproduction without permission is prohibited. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording or any information and retrieval system without permission in writing from the publisher.

All articles published, including editorial, letters and book reviews, represent the opinion of the authors and do not reflect the official policy of the publisher or the institution with which the author is affiliated, unless it is clearly specified.